Reading and Understanding Tables from the 2014 Kenya Demographic and Health Survey (KDHS)

Example 1: Exposure to Mass Media
A Question Asked of All Survey Respondents

Percentage of women age 15-49 v	vho are e	exposed to specific r	nedia on a weekly	basis, by background	l characteristics, Ke	nya 2014	
3	2	Reads a newspaper at		Listens to the radio at least once	Accesses all three media at least	Accesses none of the three media at least once a	Number of
Background characteristic		least once a week	once a week	a week	once a week	week	women
ge		04.7	22.0		0.5	05.0	- 000
15-19		21.7	33.2	66.0	9.5	25.2	5,820
20-24		19.4	44.2	73.5	12.2	18.6	5,735
25-29		18.2	45.2	71.6	11.8	20.3	6,100
30-34		16.8	41.1	70.5	10.9	22.5	4,510
35-39		14.9	36.8	68.1	9.5	25.2	3,773
40-44		15.0	31.4	67.4	9.5	28.1	2,885
45-49		15.1	32.0	68.5	9.3	25.7	2,257
esidence							
Urban		24.9	66.4	75.6	17.9	11.2	12,690
Rural		13.2	19.9	65.6	5.7	31.0	18,389
agion							
egion Coast		13.6	38.2	50.2	9.0	37.8	3,076
North Eastern		4.0	11.4	20.9		72.4	,
Eastern		17.4	24.7	67.0	1.5 8.1	29.0	648
		25.5	24.7 57.1			9.0	4,375 3,994
Central		25.5 19.1	34.6	84.1 70.3	18.1 10.9	23.4	
Rift Valley							7,953
Western		14.1	21.2	73.3	5.9	22.8	3,225
Nyanza		14.9	25.3	68.6	6.7	25.7	4,038
Nairobi		20.7	80.3	78.7	16.5	6.1	3,770
ducation							
No education		0.3	9.7	28.0	0.2	68.1	2,176
Primary incomplete		5.8	17.1	61.9	1.9	34.4	7,989
Primary complete		10.3	36.6	73.0	5.0	20.5	7,637
Secondary +		32.6	58.1	79.3	20.9	10.1	13,277
/ealth quintile							
Lowest		4.1	3.2	37.6	0.7	61.1	4,838
Second		9.9	7.5	63.3	2.2	34.2	5,457
Middle		14.6	16.5	74.6	4.3	21.5	6,032
Fourth		19.6	48.3	80.9	11.0	12.1	6,550
Highest		32.7	89.8	80.4	26.5	2.7	8,203
		4	55.5		20.5		5,203
Total		4 17.9	38.9	69.7	10.7	22.9	31,079

Step I: Read the title and subtitle. They tell you the topic and the specific population group being described. In this case, the table is about women age 15-49 and their access to different types of media. All eligible female respondents age 15-49 were asked these questions.

Step 2: Scan the column headings—highlighted in green in the table above. They describe how the information is categorized. In this table, the first three columns of data show different types of media that women access at least once a week. The fourth column shows women who access all three media, while the fifth column is women who do not access any of the three types of media at least once a week. The last column lists the number of women interviewed in the survey.

Step 3: Scan the row headings—the first vertical column highlighted in blue in the table above. These show the different ways the data are divided into categories based on population characteristics. In this case, the table presents women's access to media by age, urban-rural residence, region, educational level, and wealth quintile. Most of the tables in the KDHS report will be divided into these same categories.

Step 4: Look at the row at the bottom of the table highlighted in red. These percentages represent the totals of all women age 15-49 and their access to different types of media. In this case, 17.9% of women age 15-49 read a newspaper at least once a week, 38.9% watch television weekly, and 69.7% listen to the radio weekly.

Step 5: To find out what percentage of women with secondary+ education access all three media weekly, draw two imaginary lines, as shown on the table. This shows that 20.9% of women age 15-49 with secondary+ education access all three types of media weekly.

Practice: Use the table to the left to answer the following questions (answers are upside down, below):

- a) What percentage of women in Kenya do not access any of the three media at least once a week?
- b) What age group of women are most likely to listen to the radio weekly?
- c) Compare women in urban areas to women in rural areas—which group is more likely to watch television weekly?

Example 2: Prevalence of Diarrhoea Comparing and Understanding Patterns

Step I: Read the title and subtitle. In this case, the table presents diarrhoea among children under five in the two weeks before the survey.

Step 2: Identify the information presented in the table—highlighted in green in the table to the right. In this table there is only one indicator—diarrhoea, but it is divided into two categories: all diarrhoea and diarrhoea with blood.

Step 3: Look at the row headings to identify the background characteristics. In this table, diarrhoea is presented by age in months, sex, source of drinking water, toilet facility, urban-rural residence, region, mother's educational level, and wealth quintile.

Step 4: Look at the row in the bottom of the table to determine the total proportion of children under five with diarrhoea in the two weeks before the survey. This shows that 15.2% of children under five in Kenya had diarrhoea in the two weeks before the survey.

Step 5: In Kenya, I5% of children under five had diarrhoea in the two weeks before the survey, but a closer look at the table shows how diarrhoea varies throughout Kenya. To gain a better understanding of differences in the prevalence of diarrhoea, consider the following questions:

- Is diarrhoea more common in urban or rural areas?
 Diarrhoea is slightly more common in rural areas (15.7%) than in urban areas (14.3%). However, the difference between these two groups is small.
- Now, compare diarrhoea among girls and boys.
 Diarrhoea is slightly higher among boys (15.9%) than among girls (14.4%). However, the difference between these two groups is small.
- What are the lowest and the highest percentages (range) of diarrhoea by region? Just 7.8% of children under 5 in North Eastern had diarrhoea, compared to a high of 20.1% in Western region.
- Look for patterns: Does diarrhoea vary by background characteristics? For example, is there a clear pattern of diarrhoea by age in months? By source of drinking water? By toilet facility? By mother's education? By wealth quintile?
- Answers: Diarrhoea is highest among children age 6-11 months (26.6%), while diarrhoea is lowest among children age 48-59 months (6.7%). Children

Table 10.7 Prevalence of diarrhoea

Percentage of children under age five who had diarrhoea in the two weeks preceding the survey, by background characteristics, Kenya 2014

3	2	Diarrhoea in the preceding to	he survey	Number
Background characteristic		All diarrhoea	Diarrhoea with blood	Number of children
Age in months				
<6 6-11		12.9 26.6	1.3 2.2	1,694
12-23		26.6	2.2 1.9	1,909 3,777
24-35		15.8	2.3	3,760
36-47 48-59		9.2 6.7	1.5 1.0	3,889 3,672
		0.7	1.0	0,072
Sex Male		15.9	1.6	9,477
Female		14.4	1.8	9,477
0	.1			
Source of drinking water Improved	'	14.8	1.5	12,024
Not improved		16.1	2.2	6,496
Other/missing		7.1	0.0	182
Toilet facility ²				
Improved, not shared		11.2	1.0	3,983
Shared ³ Non-improved		16.2 16.4	1.2 2.2	4,854 9,855
·		10.1		0,000
Residence Urban		14.3	1.2	6,677
Rural		15.7	2.0	12,025
Region				
Coast		17.6	2.6	1,936
North Eastern		7.8	1.3	625
Eastern Central		14.3 10.4	0.7 0.4	2,235 1,725
Rift Valley		13.2	1.8	5,457
Western		20.1	2.7	2,166
Nyanza Nairobi		18.9 15.6	2.7 0.7	2,638 1,920
		10.0	0.7	1,020
Mother's education No education		14.1	2.8	2,218
Primary incomplete		18.4	2.4	5,304
Primary complete		14.2	1.1	5,164
Secondary+		13.7	1.2	6,016
Wealth quintile				
Lowest		17.2	2.7	4,457
Second Middle		17.1 15.5	2.3 1.7	3,803 3,375
Fourth		15.4	1.1	3,285
Highest		10.5	0.4	3,782
Total		4 15.2	1.7	18,702

Note: Total includes 14 children for whom information on toilet facility is missing.

who have access to an improved water source are less likely to have experienced diarrhoea (14.8%) than children who lack an improved source (16.1%). Children who have access to an improved, not shared toilet facility are less likely to have experienced diarrhoea (11.2%) than children who use a shared or non-improved toilet facility (16.2% and 16.4%, respectively). Diarrhoea is slightly higher among children whose mothers have no education (14.1%) than among children whose mother's have secondary+ education (13.7%). However, the difference between these two groups is very small, there is almost no difference in diarrhoea between children whose mothers have no education and children whose mothers have secondary+ education. Finally, there is also a clear pattern in diarrhoea by household wealth quintile. Diarrhoea decreases as household wealth increases; 17.2% of children living in households in the lowest wealth quintile had diarrhoea in the last two weeks, compared to 10.5% of children living in households in the highest wealth quintile.

- You can also look for patterns by diarrhoea with blood. The patterns seen in all diarrhoea can be different that the patterns by diarrhoea with blood. For example, only 1.2% of children under five had diarrohea with blood, compared to 15.2% of children with all diarrhoea.
- By looking at patterns by background characteristics, we can see which groups are more in need of interventions to address
 diarrhoea. Resources are often limited; looking for patterns can help programme planners and policymakers determine how
 to most effectively use resources.

¹ See Table 2.1 for definition of categories

² See Table 2.2 for definition of categories

³ Facilities that would be considered improved if they were not shared by two or more households

Example 3: Payment for Sexual Intercourse and Condom Use at Last Sexual Intercourse A Question Asked of a Subgroup of Survey Respondents

Table 13.10 Payment for sexual intercourse Percentage of men age 15-49 who ever months, and among them, the percentage characteristics, Kenya 2014	paid for sexual inter	course and percenta	ge reporting paym		
Uldiacteristics, Nerrya 2017	2 Percentage who	Among all men: Percentage who paid for sexual		Among men who p past 12 r Percentage reporting condom	
Background characteristic	ever paid for sexual intercourse	intercourse in the past 12 months	Number of men	use at last paid sexual intercourse	Number of men
Age					
15-24 15-19 20-24 25-29	4.0 2.2 6.1 10.3	1.9 1.4 2.5 2.5	4,666 2,540 2,125 2,104	63.5 (67.7) 60.5 69.3	88 36 52 53
30-39 40-49	12.8 10.9	3.4 2.4	3,268 2,024	(77.9) 4	48
Marital status Never married Married or living together Divorced/separated/widowed	5.6 10.1 20.7	2.2 2.2 7.8	5,350 6,095 618	73.0 72.3 82.5	117 136 48
Residence Urban Rural	10.6 7.1	2.3 2.6	5,300 6,762	79.8 70.4	121 179
Region Coast North Eastern	13.6 1.1	2.8 0.7	1,260 227	(59.0)	35 2
Eastern Central Rift Valley	11.0 5.3 5.6	1.2 1.8 2.5	1,825 1,564 3,050	(81.4) * 84.2	22 29 75
Western Nyanza Nairobi	10.0 9.1 10.7	3.6 4.3 2.3	1,164 1,405 1,568	(59.2) 72.2 *	42 60 36
Education No education Primary incomplete Primary complete Secondary+	7.5 9.9 9.9 7.4	2.2 3.5 2.6 1.9	345 3,071 2,734 5,913	* 65.0 82.0 79.2	8 108 70 115
Wealth quintile					
Lowest Second Middle Fourth Highest	8.0 7.5 8.1 9.3 9.6	2.9 2.4 3.3 2.1 2.1	1,691 2,145 2,370 2,959 2,897	65.6 68.2 71.8 83.2 (80.5)	49 52 78 61 60
Total 15-49	8.6	2.5 3	12,063	74.2	300 3
50-54	12.9	2.6	756	*	20
Total 15-54	8.9	2.5	12,819	72.8	320

Note: Figures in parentheses are based on 25-49 unweighted cases; an asterisk denotes a figure based on fewer than 25 unweighted cases that has been suppressed

Step I: Read the title and subtitle. In this case, the table is about two separate groups of men: all men age 15-49 (a) and men age 15-49 who paid for sexual intercourse in the past 12 months (b).

Step 2: Identify the two panels. First identify the columns that refer to all men age 15-49 (a), and then isolate the columns that refer only to those men who paid for sexual intercourse in the past 12 months (b).

Step 3: Look at the first panel. What percentage of men age 15-49 paid for sexual intercourse in the past 12 months? It's 2.5%. Now look at the second panel. How many men are there who paid for sexual intercourse in the past 12 months? It's 300 men or 2.5% of the 12,063 men age 15-49 (with rounding). The second panel is a subset of the first panel.

Step 4: Only 2.5% of men in the survey paid for sexual intercourse in the past 12 months. Once these men are further divided into the background characteristic categories, there may be too few cases for the percentages to be reliable.

- What percentage of men age 40-49 who paid for sex in the past 12 months reported using a condom at last paid sexual intercourse? 77.9%. This percentage is in parentheses because there are fewer than 50 men (unweighted) in this category. Readers should use this number with caution—it may not be accurate. (For more information on weighted and unweighted numbers, see Example 4.)
- What percentage of men with no education who paid for sex in the past 12 months reported using a condom at last paid sexual intercourse? There is no number in this cell—only an asterisk. This is because fewer than 25 men with no education (unweighted) paid for sex in the past 12 months. Results for this group are not reported. The subgroup is too small, and therefore the data are not reliable.

Note: When parentheses or asterisks are used in a table, the explanation will be noted under the table. If there are no parentheses or asterisks on a table, you can proceed with confidence that enough cases were included in all categories that the data are reliable.





Example 4: Understanding Sampling Weights in KDHS Tables

A sample is a group of people who have been selected for a survey. In KDHS surveys, the sample is designed to represent the national population of age 15-49. In addition to national data, most countries want to collect and report data on smaller geographical or administrative areas. However, doing so requires a minimum sample size per area (e.g., about 800 women per area). For the 2014 KDHS, the survey sample is representative of the country as a whole, for urban and rural areas, for each of Kenya's 8 regions, and for each of Kenya's 47 counties.

To generate statistics that are representative of the country as a whole and the 8 regions, the number of women surveyed in each region should contribute to the size of the total (national) sample in proportion to size of the region. However, if some regions have small populations, then a sample allocated in proportion to each region's population may not include sufficient women from each region for analysis. To solve this problem, regions with small populations are oversampled. For example, let's say that you have enough money to interview 31,079 women and want to produce results that are representative of Kenya as a whole and its regions (as in Table 3.1). However, the total population of Kenya is not evenly distributed among the regions: some regions, such as Rift Valley, are heavily populated while others, such as North Eastern are not. Thus, North Eastern must be oversampled.

A sampling statistician determines how many women should be interviewed in each region in order to get reliable statistics. The blue column (I) in the table at the right shows the actual number of women interviewed in each region. Within the regions, the number of women interviewed ranges from 999 in Nairobi region to 9,059 in Rift Valley region. The number of interviews is sufficient to get reliable results in each region.

With this distribution of interviews, some regions are overrepresented and some regions are underrepresented. For example, the population in the North Eastern region is about 2% of the population in Kenya, while Rift Valley is about 26% of the population in Kenya. But as the blue column shows, the number of women interviewed in North Eastern region accounts for about 5% of the total sample of women interviewed

Table 3.1 Background characteristics of respondents					
Percent distribution of women age 15-49 by selected background characteristics, Kenya 2014					
Background characteristic	Weighted percent	Weighted number	Unweighted number		
Region	3	2	1		
Coast	9.9	3,076	3,902		
North Eastern	2.1	648	1,664		
Eastern	14.1	4,375	5,247		
Central	12.9	3,994	3,114		
Rift Valley	25.6	7,953	9,059		
Western	10.4	3,225	2,840		
Nyanza	13.0	4,038	4,254		
Nairobi	12.1	3,770	999		
Total 15-49	100.0	31,079	31,079		

(1,664/31,079) and the number of women interviewed in Rift Valley accounts for 29% of the total sample of women interviewed (9,059/31,079). This unweighted distribution of Kenyan women does not accurately represent the population.

In order to get statistics that are representative of Kenya, the distribution of the women in the sample needs to be weighted (or mathematically adjusted) such that it resembles the true distribution in the country. Women from a small region, like North Eastern, should only contribute a small amount to the national total. Women from a large region, like Rift Valley should contribute much more. Therefore, DHS statisticians mathematically calculate a "weight" which is used to adjust the number of women from each region so that each region's contribution to the total is proportional to the actual population of the region. The numbers in the purple column (2) represent the "weighted" values. The weighted values can be smaller or larger than the unweighted values at regional level. The total national sample size of 31,079 women has not changed after weighting, but the distribution of the women in the regions has been changed to represent their contribution to the total population size.

How do statisticians weight each category? They take into account the probability that a woman was selected in the sample. If you were to compare the red column (3) to the actual population distribution of Kenya, you would see that women in each region are contributing to the total sample with the same weight that they contribute to the population of Kenya. The weighted number of women in the survey now accurately represents the proportion of women who live in North Eastern and the proportion of women who live in Rift Valley.

With sampling and weighting, it is possible to interview enough women to provide reliable statistics at national and regional levels. In general, only the weighted numbers are shown in each of the KDHS tables, so don't be surprised if these numbers seem low: they may actually represent a larger number of women interviewed.





















