

Table j: WSC Goals and Kenya's Achievements

No.	MICS INDICATOR	GOAL	ACHIEVED	
1a	Under 5 mortality	Reduction of the 1990 infant and under five mortality rate by one third or to 50 and 70 per 1,000 live births.	NO	
1b	Infant mortality		NO	
2	Maternal mortality		NOT TESTED	
3ai	Underweight prevalence (2SD)	Reduction of 1990 levels of severe and moderate malnutrition by one fifth or more.	NO	
3aii	Underweight prevalence (3SD)			
3bi	Stunting prevalence (2SD)			
3bii	Stunting prevalence (3SD)			
3ci	Wasting prevalence (2SD)			
3cii	Wasting prevalence (3SD)			
4	Use of safe drinking water	Universal access to safe drinking water.	NO/IMPROVED	
5	Sanitary means of excreta disposal	Universal sanitary means of excreta disposal.	NO	
6a	Children reaching Grade 5	Universal access to basic education, and achievement of primary education by at least 80 percent of primary school-age children, reducing disparities between boys and girls.	NO	
6b	Net primary school attendance			
6c	Proportion entering school at right age			
7	Literacy	Reduction of adult illiteracy rate to at least half its 1990 level, with emphasis on female literacy.	YES	
8	Total child disability		NOT TESTED	
9ai	Under 5 mortality (female)	Special attention to the health and nutrition of the female child and to pregnant and lactating women.	YES	
9aii	Under 5 mortality (male)		YES	
9bi	Underweight prevalence (2SD) (female)		YES	
9bii	Underweight prevalence (2SD) (male)		YES	
10a	Contraceptive prevalence (partner)		NOT TESTED	
10b	Contraceptive prevalence (all)	NOT TESTED		
11a	Antenatal care	Access by all pregnant women to prenatal care, trained attendants during childbirth and referral facilities for high-risk pregnancies and obstetric emergencies.	YES	
11b	Child birth care		NO	
12	Birth weight below 2.5 kg	Reduction of the low birth-weight rate to less than 10 per cent	YES	
14	Iodised salt	Virtual elimination of iodine deficiency disorders.	NO	
15a	Children receiving Vit A supplement	Virtual elimination of vitamin A deficiency disorders.	NO	
15b	Mothers receiving Vit A supplement		NO	
15c	Children night blind		NOT TESTED	
15d	Pregnant women night blind		NO	
16a	Exclusively breast fed	Empowerment of women to breast-feed their children exclusively for 4 to 6 months, and continue breast-feeding, with complementary food, well into the second year.	NO	
16b	Timely complementary feeding		NO	
16ci	Continued breast feeding 12-15 months		NO	
16cii	Continued breast feeding 20-23 months		NO	
22a	DPT immunisation		Maintenance of high level of immunisation coverage (at least 90 percent of children under one year of age by the year 2000) against diphtheria, pertussis, tetanus, polio, TB and against tetanus for women of child bearing age.	NO
22b	Measles immunisation			NO
22c	Polio immunisation	NO		
22d	TB immunisation	NO		
22e	Protection against neo-natal tetanus	Maintenance of high level of immunisation coverage (at least 90 percent of children under one year of age by the year 2000) in the diarrhoea incidence rate.	NO	
23a	Incidence of child diarrhea	in the diarrhoea incidence rate.	NOT TESTED	
23b	ORT		YES	
23c	Correct home management of diarrhea		NO	
24	Seek care for ARI	Reduction by one third in the deaths due to acute respiratory tract infections in children under five years.	NO TESTED	
26	Pre-school development	Expansion of early children development activities, including appropriate low-cost family - and community-based interventions.	NO	

Table 1: Number of households and women, and response rates, Kenya, 2000

	Rural/Urban		Total
	Rural	Urban	
Sampled households	7,266	1,988	9,313
Occupied households	7,251	1,863	9,140
Completed households	7,151	1,823	8,993
Household response rate	98.6	97.9	98.4
Eligible women	8,829	2,420	11,275
Interviewed women	8,019	2,243	10,281
Women response rate	90.8	92.7	91.2
Children under 5	7,046	1,655	8,726
Interviewed children under 5	5,814	1,435	7,266
Child response rate	82.5	86.7	83.3

Table 2: Single year age distribution of household population by sex, Kenya, 2000

Age	Sex				Age	Sex			
	Male		Female			Male		Female	
	Number	Percent	Number	Percent		Number	Percent	Number	Percent
0	899	4.2	857	3.6	49	75	0.3	70	0.3
1	831	3.9	828	3.5	50	198	0.9	223	0.9
2	858	4.0	806	3.4	51	89	0.4	103	0.4
3	822	3.8	831	3.5	52	154	0.7	150	0.6
4	768	3.6	748	3.2	53	64	0.3	81	0.3
5	694	3.2	662	2.8	54	75	0.4	84	0.4
6	715	3.3	679	2.9	55	128	0.6	112	0.5
7	677	3.1	683	2.9	56	74	0.3	59	0.3
8	739	3.4	748	3.2	57	67	0.3	55	0.2
9	617	2.9	664	2.8	58	64	0.3	61	0.3
10	697	3.2	726	3.1	59	39	0.2	32	0.1
11	614	2.9	606	2.6	60	96	0.4	117	0.5
12	696	3.2	671	2.9	61	67	0.3	63	0.3
13	620	2.9	627	2.7	62	60	0.3	42	0.2
14	683	3.2	712	3.0	63	40	0.2	24	0.1
15	516	2.4	512	2.2	64	50	0.2	45	0.2
16	549	2.6	563	2.4	65	71	0.3	70	0.3
17	400	1.9	473	2.0	66	39	0.2	30	0.1
18	472	2.2	558	2.4	67	50	0.2	16	0.1
19	257	1.2	361	1.5	68	36	0.2	43	0.2
20	390	1.8	536	2.3	69	18	0.1	20	0.1
21	249	1.2	407	1.7	70+	307	1.4	252	1.1
22	265	1.2	496	2.1	Missing/C	3	0.0	6	0.0
23	238	1.1	366	1.6		21,516	100	23,488	100
24	272	1.3	412	1.8					
25	270	1.3	441	1.9					
26	276	1.3	430	1.8					
27	242	1.1	399	1.7					
28	270	1.3	423	1.8					
29	193	0.9	246	1.0					
30	385	1.8	426	1.8					
31	221	1.0	279	1.2					
32	309	1.4	347	1.5					
33	175	0.8	184	0.8					
34	177	0.8	245	1.0					
35	294	1.4	313	1.3					
36	233	1.1	270	1.2					
37	200	0.9	224	1.0					
38	239	1.1	280	1.2					
39	147	0.7	187	0.8					
40	290	1.3	316	1.3					
41	151	0.7	165	0.7					
42	205	1.0	198	0.8					
43	112	0.5	138	0.6					
44	111	0.5	160	0.7					
45	166	0.8	124	0.5					
46	141	0.7	134	0.6					
47	122	0.6	103	0.4					
48	185	0.9	170	0.7					

Table 3: Percentage of cases missing information for selected questions , Kenya, 2000

	Percent missing	Number
Level of education	26.7	37,292
Year of education	8.4	37,292
Number of hours worked for pay	90.0	4,767
Complete birth date	35.7	10,281
Date of last tetanus toxoid injection	63.3	1,621
Ever been tested for HIV	7.6	10,278
Complete birth date	6.7	7,266
Diarrhoea in last 2 weeks	1.7	7,266
Weight	7.8	7,266
Height	11.8	7,266

Table 4: Percent distribution of households by background characteristics, Kenya, 2000

	Rural/Urban		Total
	Rural	Urban	
Province			
Nairobi	0.0	57.5	14.2
Central	17.1	4.4	14.0
Coast	6.4	12.5	7.9
Eastern	18.7	3.6	15.0
North Eastern(Urban)	0.0	3.7	0.9
Nyanza	20.3	4.2	16.3
Rift valley	23.7	12.3	20.9
Western	13.8	1.8	10.8
Number of HH members			
1	3.5	3.5	3.5
2-3	20.6	30.1	22.9
4-5	36.1	41.3	37.4
6-7	24.7	17.9	23.0
8-9	10.9	5.3	9.5
10+	4.3	1.9	3.7
Total	100	100	100
At least one child age < 15	89.4	90.5	89.6
At least one child age < 5	62.1	68.5	63.7
At least one woman age 15-49	86.5	94.1	88.4
Number	6,747	2,218	8,965
weighted	7,149	1,822	8,971

Table 5: Percent distribution of women 15-49 by background characteristics, Kenya, 2000

	Percent	Number	Unweighted
Province			
Nairobi	16.1	1,656	486
Central	12.6	1,293	1,352
Coast	8.8	909	1,319
Eastern	15.2	1,566	1,371
North Eastern(Urban)	1.0	98	103
Nyanza	15.6	1,607	1,752
Rift Valley	20.2	2,073	2,919
Western	10.5	1,079	979
Rural/Urban			
Rural	72.9	7,492	8,035
Urban	27.1	2,789	2,246
Age			
15-19	21.2	2,179	2,196
20-24	20.4	2,097	2,032
25-29	18.2	1,870	1,806
30-34	13.7	1,413	1,461
35-39	12.0	1,234	1,261
40-44	9.0	927	949
45-49	5.5	561	576
Ever given birth			
Ever gave birth	75.5	7,763	7,735
Never gave birth	24.2	2,483	2,520
Missing	0.3	35	26
Woman's education level			
None	15.3	1,578	1,733
Primary	57.2	5,882	5,956
Secondary +	27.4	2,821	2,592
Total	100	10,281	10,281

Table 6: Percent distribution of children under 5 by background characteristics, Kenya, 2000

	Percent	Number	Unweighted
Sex			
Male	51.2	3,720	3,724
Female	48.8	3,546	3,542
Province			
Nairobi	15.5	1,130	321
Central	10.9	794	842
Coast	9.5	689	938
Eastern	10.6	768	667
North Eastern(Urban)	1.1	83	83
Nyanza	16.0	1,161	1,179
Rift valley	24.4	1,775	2,462
Western	11.9	867	774
Rural/Urban			
Rural	73.7	5,357	5,827
Urban	26.3	1,909	1,439
Age			
< 6 months	10.6	768	721
6-11 months	10.8	781	763
12-23 months	21.2	1,544	1,530
24-35 months	20.2	1,470	1,458
36-47 months	19.7	1,430	1,462
48-59 months	17.5	1,272	1,332
Woman's education level			
None	22.5	1,637	1,750
Primary	53.2	3,864	3,887
Secondary +	24.3	1,765	1,629
Total	100	7,266	7,266

Table 9: Percentage of children aged 36-59 months who are attending some form of organized early childhood education programme

	Attending programme	Number of children
Sex		
Male	14.1	1,405
Female	17.3	1,297
Province		
Nairobi	30.9	368
Central	18.9	319
Coast	14.2	245
Eastern	10.1	323
North Eastern(Urban)	12.7	28
Nyanza	12.8	414
Rift valley	13.9	691
Western	9.0	314
Rural/Urban		
Rural	11.0	2,049
Urban	30.1	654
Age		
36-47 months	8.6	1,430
48-59 months	23.5	1,272
Woman's education level		
None	12.4	704
Primary	10.7	1,354
Secondary +	29.5	644
Total	15.6	2,702

World Summit for Children Goal => Number 26

Table 10: Percentage of children entering first grade of primary school who eventually reach grade 5, Kenya, 2000

	Percent in grade 1 reaching grade 2	Percent in grade 2 reaching grade 3	Percent in grade 3 reaching grade 4	Percent in grade 4 reaching grade 5	Percent who reach grade 5 of those who enter grade 1
Sex					
Male	96.7	97.4	97.0	95.9	87.5
Female	98.4	97.7	97.6	96.4	90.4
Province					
Nairobi	100.0	100.0	97.4	100.0	97.4
Central	97.9	96.4	98.5	97.6	90.7
Coast	96.4	95.9	97.2	90.8	81.6
Eastern	96.8	97.7	98.5	96.0	89.5
North Eastern(Urban)	100.0	100.0	100.0	90.2	90.2
Nyanza	96.3	97.9	97.7	95.5	88.0
Rift valley	99.0	97.7	97.6	96.9	91.6
Western	95.7	96.8	93.7	94.6	82.1
Rural/Urban					
Rural	97.3	97.5	97.3	96.1	88.7
Urban	98.7	97.5	97.0	96.4	90.0
TOTAL	97.5	97.5	97.3	96.1	88.9

World Summit for Children Goal => Number 6

Table 10a: Percentage of children entering first Std of primary school who eventually reach std 8, Kenya, 2000

	Percent in Std 1 reaching Std 2	Percent in Std 2 reaching Std 3	Percent in Std 3 reaching Std 4	Percent in Std 4 reaching Std 5	Percent in Std 5 reaching Std 6	Percent in Std 6 reaching Std 7	Percent in Std 7 reaching Std 8	Percent who reach Std 8 of those who enter Std 1
Sex								
Male	96.7	97.4	97.0	95.9	95.9	95.6	95.2	76.4
Female	98.4	97.7	97.6	96.4	97.1	95.7	95.7	80.4
Province								
Nairobi	100.0	100.0	97.4	100.0	96.2	100.0	100.0	93.7
Central	97.9	96.4	98.5	97.6	98.5	96.5	93.8	80.9
Coast	96.4	95.9	97.2	90.8	100.0	95.7	93.5	73.0
Eastern	96.8	97.7	98.5	96.0	99.0	99.4	100.0	88.0
North Eastern (Urban)	100.0	100.0	100.0	90.2	100.0	88.5	82.9	66.2
Nyanza	96.3	97.9	97.7	95.5	93.8	91.7	96.4	73.0
Rift valley	99.0	97.7	97.6	96.9	95.0	96.4	92.2	77.3
Western	95.7	96.8	93.7	94.6	95.9	93.1	96.2	70.5
Rural/Urban								
Rural	97.3	97.5	97.3	96.1	96.5	95.2	95.6	77.9
Urban	98.7	97.5	97.0	96.4	96.5	97.8	95.0	80.6
Wealth Index Quintiles								
Poorest	95.7	98.4	97.2	92.1	92.3	91.5	95.9	68.2
Second	96.1	97.4	97.3	95.6	98.6	94.5	92.3	74.8
Middle	98.2	97.5	96.7	98.2	95.2	97.2	96.1	80.9
Fourth	99.2	96.7	97.0	97.7	98.7	96.6	97.6	84.5
Richest	100.0	97.4	98.9	98.7	97.8	98.5	95.5	87.4
Total	97.5	97.5	97.3	96.1	96.5	95.7	95.5	78.4

World Summit for Children Goal => Number 6

Table 11: Percentage of children of primary school age attending primary school, Kenya, 2000

	Sex					
	Male		Female		Total	
	Attending primary school		Attending primary school			
	Attending	Number	Attending	Number	Attending	Number
Province						
Nairobi	68.5	486	80.4	448	74.2	934
Central	82.5	648	83.7	599	83.1	1,247
Coast	59.9	495	53.8	456	57.0	951
Eastern	71.2	874	78.0	959	74.8	1,833
North Eastern(Urban)	61.6	35	58.7	45	60.0	80
Nyanza	75.8	893	74.4	925	75.1	1,817
Rift valley	73.3	1,247	75.2	1,297	74.2	2,545
Western	72.9	696	73.6	674	73.2	1,371
Rural/Urban						
Rural	72.7	4,461	74.3	4,509	73.5	8,971
Urban	72.4	913	77.3	894	74.8	1,807
Age						
6	29.5	715	33.7	679	31.6	1,394
7	57.3	677	57.5	683	57.4	1,360
8	72.0	739	76.8	748	74.4	1,486
9	83.6	617	82.6	664	83.0	1,280
10	84.1	697	87.4	726	85.8	1,424
11	86.6	614	89.0	606	87.8	1,220
12	86.9	696	87.5	671	87.2	1,366
13	86.8	620	85.5	627	86.2	1,247
Total	72.7	5,374	74.8	5,403	73.7	10,777

World Summit for Children Goal => Number 6

Table 12: Percentage of the population aged 15 years and older that is literate, Kenya, 2000

	Sex						Literate	Total Not known	Number
	Male			Female					
	Literate	Not known	Number	Literate	Not known	Number			
Province									
Nairobi	84.0	0.0	1,457	81.1	0.0	1,886	82.4	0.0	3,343
Central	86.5	0.0	1,375	81.9	0.0	1,647	84.0	0.0	3,022
Coast	72.1	0.0	882	55.7	0.0	1,088	63.0	0.0	1,970
Eastern	68.7	0.0	1,659	64.6	0.0	1,975	66.5	0.0	3,634
North Eastern(Urban)	76.9	0.0	101	51.3	0.0	99	64.2	0.0	200
Nyanza	76.2	0.0	1,675	66.4	0.0	2,023	70.9	0.0	3,699
Rift valley	76.6	0.0	2,221	69.1	0.0	2,522	72.6	0.0	4,743
Western	80.2	0.0	1,214	69.7	0.0	1,395	74.6	0.0	2,609
Rural/Urban									
Rural	75.0	0.0	8,079	66.4	0.0	9,495	70.4	0.0	17,575
Urban	86.3	0.0	2,504	81.6	0.0	3,141	83.7	0.0	5,645
Age									
15-24	79.8	0.0	3,608	80.7	0.0	4,684	80.3	0.0	8,292
25-34	86.2	0.0	2,518	81.4	0.0	3,418	83.4	0.0	5,936
35-44	80.5	0.0	1,982	65.5	0.0	2,250	72.5	0.0	4,233
45-54	74.0	0.0	1,269	46.5	0.0	1,242	60.4	0.0	2,512
55-64	58.9	0.0	686	28.1	0.0	610	44.4	0.0	1,296
65+	44.0	0.0	521	20.0	0.0	432	33.1	0.0	952
Total	77.6	0.0	10,584	70.2	0.0	12,636	73.6	0.0	23,220

World Summit for Children Goal => Number 7

Table 13: Percentage of the population using improved drinking water sources, 2000

	Main source of water												Total with safe drinking water	Number of persons
	Piped into dwelling	Piped into yard or plot	Public tap	Tubewell/borehole with pump	Protected dug well	Protected spring	Rainwater collection	Bottled water	Unprotected dug well/Unprotected spring	Pond, river or stream	Tanker truck vendor	Other		
Province														
Nairobi	31.7	30.7	31.4	0.2	0.0	0.0	0.0	0.0	0.0	0.7	3.9	1.2	93.9	5,691
Central	15.3	14.1	3.6	3.2	6.0	5.8	1.9	0.1	7.1	41.8	0.7	0.4	49.7	5,541
Coast	11.9	2.6	36.3	10.2	2.3	0.2	3.9	0.0	5.4	20.6	5.8	0.7	67.1	3,936
Eastern	5.8	10.0	11.3	5.0	2.9	1.4	0.7	0.1	20.4	38.8	1.5	2.0	37.2	7,133
North Eastern (Urban)	9.0	49.7	1.6	3.4	10.9	0.0	0.0	0.0	8.1	1.6	6.4	9.3	74.6	395
Nyanza	3.7	1.8	5.9	12.1	2.8	15.7	1.9	0.0	16.6	37.8	1.2	0.5	44.0	7,344
Rift valley	9.8	8.2	7.3	4.4	12.9	5.7	1.6	0.1	11.5	34.4	2.5	1.7	49.8	9,898
Western	6.1	3.9	5.0	15.6	8.5	11.1	0.4	0.1	33.1	14.6	0.6	0.9	50.6	5,222
Rural/Urban														
Rural	6.0	5.9	6.5	8.6	7.2	7.8	1.7	0.1	17.3	37.2	0.8	1.1	43.5	35,175
Urban	29.8	26.4	31.1	0.6	1.0	0.5	0.5	0.0	0.8	1.4	6.4	1.5	89.7	9,983
Wealth Index Quintiles														
Poorest	0.0	0.0	0.8	1.7	1.0	1.8	0.3	0.0	4.2	9.9	0.1	0.2	28.0	9,800
Second	0.3	0.1	1.1	1.8	1.9	2.1	0.2	0.0	4.4	7.8	0.2	0.2	37.7	9,550
Middle	1.3	2.0	2.7	1.5	1.4	1.2	0.2	0.0	3.0	6.4	0.2	0.2	50.6	9,155
Fourth	1.9	2.6	4.3	1.4	1.3	0.9	0.4	0.0	1.5	4.0	0.9	0.5	63.8	8,671
Richest	8.3	6.2	3.6	0.2	0.2	0.0	0.2	0.0	0.1	0.2	0.8	0.1	93.7	7,982
Total	11.9	10.9	12.6	6.6	5.6	6.0	1.4	0.1	13.2	28.4	2.2	1.2	54.8	45,159

World Summit for Children Goal => Number 4

Table 14: Percentage of the population using sanitary means of excreta disposal, 2000

	Type of toilet facility								Total	Total with sanitary means of excreta disposal	Number of persons
	Flush to sewage system/septic tank	Pour flush latrine	Improved pit latrine	Traditional pit latrine	Open pit	Bucket	Other	No facilities/brush/field			
Province											
Nairobi	42.1	6.0	6.4	41.0	2.5	0.5	0.0	1.5	100	95.4	5,691
Central	10.6	0.3	6.9	80.3	1.8	0.0	0.1	0.0	100	97.8	5,541
Coast	12.3	2.4	4.4	49.3	4.6	0.0	0.1	26.9	100	68.0	3,936
Eastern	7.5	0.1	9.2	59.7	10.2	0.0	0.1	13.3	100	76.4	7,133
North Eastern(Urban)	25.3	1.6	3.1	49.2	10.9	8.4	1.6	0.0	100	79.2	395
Nyanza	5.8	0.6	4.3	59.2	3.5	0.0	0.1	26.5	100	69.9	7,344
Rift valley	10.5	0.7	7.7	53.3	8.0	0.2	0.3	19.3	100	71.9	9,803
Western	9.3	0.7	5.1	76.5	3.3	0.0	0.0	5.1	100	91.5	5,317
Rural/Urban											
Rural	6.0	0.4	6.4	64.0	6.0	0.0	0.1	17.2	100	76.6	35,175
Urban	38.1	4.7	6.9	45.4	2.6	0.8	0.3	1.3	100	94.8	9,983
Total	13.9	1.4	6.5	59.4	5.2	0.2	0.1	13.2	100	81.1	45,159

World Summit for Children Goal => Number 5

Table 15: Percentage of under-five children who are severely or moderately undernourished, 2000

	Weight for age: -2 SD	Weight for age: -3 SD	Height for age: -2 SD	Height for age: -3 SD	Weight for height: - 2 SD	Weight for height: - 3 SD	Number of children
Sex							
Male	22.6	6.7	37.9	15.7	6.6	1.8	3,000
Female	19.6	4.6	32.6	13.7	5.3	0.9	2,917
Province							
Nairobi	12.4	1.8	29.6	11.1	3.1	0.8	760
Central	15.4	3.3	27.4	9.6	4.6	1.1	627
Coast	21.1	4.0	33.7	15.5	6.4	1.5	566
Eastern	29.6	6.9	42.8	17.4	7.8	1.4	673
North Eastern(Urban)	16.6	3.7	35.4	17.0	7.2	0.0	65
Nyanza	19.9	6.2	35.9	16.3	5.2	1.7	959
Rift valley	24.9	7.3	36.8	14.9	7.6	1.5	1,533
Western	21.5	8.0	38.1	16.9	5.5	1.4	734
Rural/Urban							
Rural	23.9	6.8	38.0	16.0	6.8	1.5	4,497
Urban	12.4	2.1	26.6	10.4	3.3	0.8	1,420
Age							
< 6 months	3.0	0.5	12.4	1.4	2.4	0.8	322
6-11 months	14.6	4.2	24.5	6.2	3.9	0.6	598
12-23 months	28.4	9.2	47.5	21.3	9.9	2.5	1,312
24-35 months	22.5	6.6	34.8	13.5	6.6	1.4	1,290
36-47 months	19.4	3.8	34.5	15.3	4.1	1.0	1,286
48-59 months	21.9	4.9	34.7	16.1	4.9	0.9	1,109
Woman's education level							
None	24.1	7.2	37.2	17.2	7.1	1.6	1,347
Primary	23.3	6.8	38.8	16.3	6.6	1.7	3,147
Secondary +	13.7	1.8	25.6	8.9	3.4	0.5	1,423
Wealth Index Quintiles							
Poorest	29.1	9.3	45.3	20.9	7.6	2.4	1,390
Second	23.0	6.3	38.4	14.4	7.0	1.1	1,309
Middle	25.4	6.7	37.9	17.5	6.6	1.4	1,083
Fourth	16.4	3.7	29.0	10.9	4.4	0.6	1,090
Richest	8.6	1.2	22.0	7.9	3.5	1.0	1,045
Total	21.2	5.7	35.3	14.7	6.0	1.4	5,917

World Summit for Children Goal => Number 3, 9

Table 16: Percent of living children by breastfeeding status, Kenya, 2000

	Exclusive breastfeeding		Complementary feeding rate		Continued breastfeeding rate		Continued breastfeeding rate	
	Children 0-3 months	Number of children	Children 6-9 months	Number of children	Children 12-15 months	Number of children	Children 20-23 months	Number of children
Sex								
Male	17.9	274	71.0	255	83.0	301	41.5	210
Female	14.3	237	70.7	254	83.0	279	32.8	237
Province								
Nairobi	28.5	100	78.4	104	67.5	88	21.3	69
Central	11.4	41	80.8	57	80.0	55	28.7	43
Coast	8.1	46	70.9	46	86.9	57	45.7	33
Eastern	14.5	32	81.7	38	92.5	55	50.2	49
North Eastern(Urban)	19.0	6	68.9	4	80.7	12	0.0	4
Nyanza	8.0	105	57.9	89	89.9	110	44.8	71
Rift valley	15.0	110	81.0	115	83.8	137	44.2	106
Western	21.7	70	38.9	55	82.3	67	27.5	73
Rural/Urban								
Rural	13.7	366	68.4	349	87.4	428	40.8	324
Urban	22.6	145	76.2	160	70.8	153	26.5	123
Woman's education level								
None	17.4	83	74.0	85	76.0	101	37.1	105
Primary	15.2	316	71.1	286	84.8	337	37.2	228
Secondary +	18.3	111	68.3	138	83.7	143	36.1	114
Total	16.2	511	70.8	509	83.0	581	36.9	447

World Summit for Children Goal => Number 16

Table 16w: Percent distribution of children by breastfeeding status, Kenya, 2000

	Breastfeeding status					Total	Number of children
	Not breastfeeding	Exclusively breastfed	Breast milk and water only	Breast milk and liquids only	Breast milk and solid/mushy food		
Age							
0-1	1.0	24.5	24.3	42.2	8.1	100	247
2-3	0.6	8.3	8.5	55.4	27.2	100	243
4-5	3.5	2.8	9.5	29.0	55.2	100	252
6-7	2.0	0.5	1.3	27.8	68.5	100	256
8-9	3.6	0.1	1.4	18.1	76.7	100	241
10-11	10.0	0.4	2.2	14.2	73.2	100	268
12-13	13.0	0.0	2.0	11.4	73.5	100	290
14-15	17.3	0.3	2.2	16.9	63.3	100	278
16-17	24.8	0.0	2.4	11.9	60.8	100	276
18-19	31.3	0.0	0.9	15.9	51.9	100	212
20-21	54.7	0.0	0.3	2.8	42.3	100	201
22-23	64.5	0.0	1.4	5.0	29.1	100	205
24-25	71.5	0.3	0.5	4.0	23.6	100	249
26-27	77.4	0.0	0.2	3.4	19.0	100	284
28-29	81.4	0.0	0.1	4.6	13.9	100	266
30-31	91.1	0.0	0.0	0.9	8.1	100	192
32-33	86.7	0.2	0.0	5.2	7.9	100	165
34-35	96.8	0.0	0.0	0.6	2.6	100	218

Exclusively breastfed includes vitamin, mineral supplements and medicine

Table 17: Percentage of population consuming adequately iodized salt, Kenya, 2000

	Percent of population with no salt	Percent of population in which salt was tested	Result of test		Number of persons
			< 15 PPM	15+ PPM	
Province					
Nairobi	3.3	95.0	15.8	84.2	5,691
Central	1.8	97.6	17.7	82.3	5,541
Coast	1.2	98.0	5.9	94.1	3,936
Eastern	2.0	97.2	11.4	88.6	7,133
North Eastern(Urban)	12.8	85.8	14.5	85.5	395
Nyanza	2.2	97.1	7.3	92.7	7,344
Rift valley	2.8	96.8	3.2	96.8	9,898
Western	3.4	96.3	3.9	96.1	5,222
Rural/Urban					
Rural	2.4	97.0	8.5	91.5	35,175
Urban	3.0	95.7	11.9	88.1	9,983
Total	2.6	96.7	9.3	90.7	45,159

World Summit for Children Goal => Number 14

Table 18: Percent distribution of children aged 6-59 months by whether they received a high dose of Vitamin A supplement in the last 6 months, 2000

	Vitamin A					Total	Number of children
	Received: within last 6 months	Received: prior to last 6 months	Received: not sure when	Not sure if received	Not received		
Sex							
Male	9.5	32.1	0.3	1.5	56.6	100	3,251
Female	10.1	29.4	0.6	1.0	58.9	100	3,132
Province							
Nairobi	6.4	23.6	0.3	0.0	69.7	100	959
Central	7.7	24.4	1.4	0.9	65.6	100	701
Coast	10.4	28.6	0.5	1.7	58.8	100	608
Eastern	6.1	25.7	0.2	0.7	67.3	100	696
North Eastern(Urban)	17.9	22.4	0.0	4.6	55.1	100	73
Nyanza	11.5	42.2	0.2	1.0	45.1	100	1,003
Rift valley	12.0	30.5	0.5	1.8	55.1	100	1,588
Western	11.4	38.3	0.0	2.4	47.8	100	755
Rural/Urban							
Rural	10.8	32.2	0.4	1.1	55.5	100	4,730
Urban	7.1	26.7	0.5	1.7	63.9	100	1,652
Age							
6-11 months	17.8	10.3	0.0	0.4	71.5	100	766
12-23 months	13.4	31.6	0.4	0.8	53.8	100	1,511
24-35 months	7.5	35.0	0.5	1.1	55.9	100	1,441
36-47 months	7.6	33.1	0.4	1.5	57.3	100	1,413
48-59 months	5.7	34.6	0.7	2.4	56.5	100	1,252
Woman's education level							
None	9.6	24.3	0.4	3.2	62.5	100	1,465
Primary	9.7	33.1	0.4	0.7	56.1	100	3,345
Secondary +	10.2	31.9	0.4	0.8	56.6	100	1,573
Total	9.8	30.8	0.4	1.3	57.7	100	6,383

Table 19: Percentage of women with a birth in the last 12 months by whether they received a high dose of Vitamin A supplement before the infant was 8 weeks old, 2000

	Received Vitamin A supplement	Not sure if received	Number of women
Province			
Nairobi	25.1	12.4	342
Central	18.3	15.7	182
Coast	30.3	12.6	159
Eastern	13.9	9.9	202
North Eastern(Urban)	8.4	87.4	19
Nyanza	26.2	9.3	335
Rift Valley	22.5	12.3	370
Western	25.3	16.5	218
Rural/Urban			
Rural	21.9	12.4	1,320
Urban	26.3	15.1	508
Woman's education level			
None	22.1	21.6	234
Primary	21.3	11.1	1,105
Secondary +	27.7	13.7	489
Total	23.1	13.2	1,828

World Summit for Children Goal => Number 15

19a: Proportion of women who had night blindness during the last pregnancy

	Per cent	No. of Women
Province		
Nairobi	10.0	90
Central	6.2	162
Coast	12.6	198
Eastern	10.7	177
North Eastern		3
Nyanza	14.4	320
Rift Valley	10.9	496
Western	14.2	176
Total	11.6	1,622
Rural/Urban		
Rural	11.7	1,315
Urban	11.1	307
Total	11.6	1,622
Age		
15-19	11.6	207
20-24	9.5	516
25-29	10.9	412
30-34	14.9	275
35-39	12.3	146
40-44	17.6	51
45-49	13.3	15
Total	11.6	1,622
Woman's education level		
None	13.7	241
Primary	11.6	1,000
Secondary +	10.2	381
Total	11.6	1,622

Table 20: Percentage of live births in the last 12 months that weighed below 2500 grams at birth, Kenya, 2000

	Percent of live births below 2500 grams	Percent of live births weighed at birth	Number of live births
Province			
Nairobi	9.9	57.8	342
Central	8.4	42.5	182
Coast	9.2	33.5	159
Eastern	10.0	46.6	202
North Eastern(Urban)	2.3	0.0	19
Nyanza	8.9	20.6	335
Rift Valley	9.4	28.2	370
Western	9.4	21.4	218
Rural/Urban			
Rural	9.2	28.0	1,320
Urban	9.5	53.9	508
Woman's education level			
None	8.4	23.2	234
Primary	9.5	29.7	1,105
Secondary +	9.3	53.2	489
Total	9.3	35.2	1,828

World Summit for Children Goal => Number 12

Table 20a. Birth weight and size at birth

	Birth weight			Size of child				Total	Number
	<2500	2500+	DK/Missing	Large/Big	Average	Small	Don't know		
Province									
Nairobi	2.8	55.0	42.2	35.4	48.4	16.2	0.0	100	304
Central	3.2	39.3	57.5	34.7	54.2	11.1	0.0	100	159
Coast	3.8	29.8	66.5	26.6	55.1	14.3	4.0	100	140
Eastern	8.6	38.0	53.4	29.7	55.2	15.0	0.0	100	184
North Eastern(Urban)	0.0	0.0	100.0	0.0	66.1	33.9	0.0	100	2
Nyanza	2.5	18.0	79.4	41.5	45.2	12.6	0.7	100	306
Rift Valley	3.0	25.2	71.8	29.3	56.2	13.6	1.0	100	332
Western	1.6	19.7	78.6	37.2	48.4	14.4	0.0	100	193
Rural/Urban									
Rural	3.3	24.7	72.0	35.0	50.7	13.6	0.7	100	1,178
Urban	3.7	50.2	46.1	31.2	52.9	15.1	0.7	100	441
Woman's education level									
None	3.3	19.9	76.8	33.1	51.6	14.8	0.5	100	184
Primary	2.9	26.8	70.3	35.5	49.1	14.4	1.0	100	994
Secondary +	4.6	48.6	46.8	31.0	56.1	12.9	0.0	100	442
Total	3.4	31.8	64.8	34.0	51.3	14.0	0.7	100	1,620

Table 21 Percentage of children 12-23 months immunized against childhood diseases at any time before the survey, Kenya, 2000

	BCG	DPT1	DPT2	DPT3	Polio 0	Polio 1	Polio 2	polio3	Measles	All	None	Number of children
Vaccinated at any time before the survey												
According to:												
Vaccination Card	63.2	63.3	61.0	58.4	55.7	62.5	61.0	57.7	51.2	47.8	0.2	1,544
Mother's Report	27.7	26.1	21.7	17.8	25.0	24.1	20.1	15.1	24.9	13.0	5.6	
Either	90.9	89.4	82.7	76.2	80.7	86.6	81.1	72.8	76.1	60.8	5.7	1,544
Vaccinated by 12 months of age	90.5	88.9	82.2	75.1	80.2	85.7	80.3	71.7	71.6	55.6	0.0	

World Summit for Children Goal => Number 22

Table 22: Percentage of children age 12-23 months currently vaccinated against childhood diseases, 2000

	BCG	DPT 1	DPT 2	DPT 3	Polio 0	Polio 1	Polio 2	Polio 3	Measles	All	None	% with health card	Number of children
Sex													
Male	90.3	88.1	82.6	77.3	79.8	84.7	79.9	73.1	76.5	61.9	6.7	64.3	770
Female	91.5	90.8	82.8	75.1	81.6	88.5	82.3	72.5	75.8	59.7	4.7	66.4	774
Province													
Nairobi	95.5	92.4	86.4	84.8	93.9	93.9	85.0	74.3	84.9	64.0	3.0	61.1	225
Central	88.8	87.1	84.5	82.7	81.7	83.3	81.1	77.6	80.2	69.0	5.9	64.9	147
Coast	93.7	91.9	81.3	78.2	84.1	91.2	84.6	75.1	76.8	61.1	4.2	74.7	150
Eastern	91.5	90.5	83.9	76.0	80.0	86.9	79.9	74.5	83.2	68.2	6.0	63.6	173
North Eastern(Urban)	95.7	90.0	78.5	72.8	90.0	90.0	84.2	84.2	90.0	72.8	4.3	74.2	21
Nyanza	85.7	83.3	72.0	61.7	72.6	75.7	69.4	56.9	68.4	47.7	10.2	57.2	257
Rift valley	93.2	91.7	87.4	80.7	86.4	89.7	86.8	80.3	79.5	68.2	4.5	70.4	378
Western	86.3	88.6	82.7	70.6	61.2	84.7	79.3	69.2	58.1	45.4	6.2	65.2	191
Rural/Urban													
Rural	89.3	88.3	81.9	74.1	76.5	84.0	79.3	71.5	73.1	58.7	6.7	65.1	1,150
Urban	95.6	92.8	85.1	82.4	93.1	94.2	86.4	76.3	85.0	66.9	2.9	66.1	394
Woman's education level													
None	85.1	82.2	73.2	66.0	71.1	80.0	71.8	63.3	68.5	53.5	10.7	59.8	318
Primary	91.7	90.3	82.8	75.6	80.0	86.2	81.4	72.2	74.8	58.2	4.9	66.1	851
Secondary +	94.0	93.6	90.5	86.2	90.6	92.9	88.3	82.1	85.7	72.8	3.4	68.4	375
Total	90.9	89.4	82.7	76.2	80.7	86.6	81.1	72.8	76.1	60.8	5.7	65.4	1,544

Table 23: Percentage of under-five children with diarrhea in the last two weeks and treatment with ORS or ORT, 2000

	Diarrhea	Number of children under 5	Breast milk	Gruel	Local acceptable	ORS packet	Other milk or infant formula	Water with feeding	Any recommend ed treatment	No treatment	Number of children with diarrhoea
Sex											
Male	17.1	3,720	41.1	42.2	21.6	39.6	31.5	52.3	75.8	24.2	637
Female	14.9	3,546	45.7	50.4	23.0	38.7	27.2	53.9	77.4	22.6	527
Province											
Nairobi	15.2	1,130	46.2	40.7	14.6	61.6	35.9	53.0	85.8	14.2	172
Central	11.4	794	47.2	54.0	30.3	51.0	36.9	50.0	79.8	20.2	90
Coast	19.3	689	44.5	33.2	26.5	33.2	15.7	63.4	73.9	26.1	133
Eastern	13.7	768	45.7	40.4	21.4	26.1	23.9	35.3	71.2	28.8	105
North Eastern(Urban)	10.5	83	24.1	13.8	24.1	27.0	13.8	16.7	27.0	73.0	9
Nyanza	19.8	1,161	41.7	55.7	29.7	34.0	33.0	57.7	78.7	21.3	230
Rift valley	14.1	1,775	42.7	49.6	20.2	35.7	36.1	53.9	75.8	24.2	250
Western	20.1	867	39.1	43.3	15.8	35.9	20.1	51.8	71.6	28.4	174
Rural/Urban											
Rural	15.8	5,357	42.3	45.7	22.5	33.8	28.0	51.3	74.3	25.7	847
Urban	16.6	1,909	45.4	46.5	21.6	53.5	33.7	57.5	82.5	17.5	317
Age											
< 6 months	12.4	768	67.5	24.5	16.0	34.7	25.1	30.8	72.0	28.0	95
6-11 months	30.7	782	72.9	41.8	15.7	45.0	34.4	46.4	79.7	20.3	240
12-23 months	22.4	1,544	57.2	49.9	27.2	39.0	30.0	58.3	76.9	23.1	345
24-35 months	14.7	1,470	21.6	56.7	18.2	43.6	28.9	60.9	79.4	20.6	216
36-47 months	10.3	1,430	9.5	39.6	26.2	31.5	26.7	48.7	68.7	31.3	147
48-59 months	9.4	1,272	4.3	47.8	28.4	33.1	26.9	59.7	77.1	22.9	120
Woman's education level											
None	16.3	1,637	37.8	36.4	19.6	30.5	24.4	52.5	72.8	27.2	266
Primary	17.4	3,864	43.8	49.9	23.1	42.1	27.5	53.2	78.0	22.0	673
Secondary +	12.7	1,765	47.8	45.0	22.9	40.8	41.7	53.2	76.4	23.6	224
Total	16.0	7,266	43.2	45.9	22.2	39.2	29.5	53.0	76.5	23.5	1,164

World Summit for Children Goal => Number 23

Table 24: Percentage of under-five children with diarrhea in the last two weeks who took increased fluids and continued to feed during the episode, 2000

	Diarrhea	Number of children under 5	Drinking during diarrhea			Total	Eating during diarrhea			Total	Received increased fluids and continued eating	Number of children with diarrhea
			More	Same/Less	Missing/DK		somewhat less/same/more	Much less/none	Missing/DK			
Sex												
Male	17.1	3,720	22.5	59.1	18.4	100	27.8	67.4	4.8	100	8.5	637
Female	14.9	3,546	20.1	63.4	16.5	100	24.4	68.0	7.6	100	5.3	527
Province												
Nairobi	15.2	1,130	18.5	75.6	5.9	100	8.4	89.4	2.2	100	1.7	172
Central	11.4	794	33.2	54.4	12.4	100	28.2	67.3	4.5	100	10.9	90
Coast	19.3	689	31.9	53.1	15.0	100	39.6	58.0	2.4	100	13.0	133
Eastern	13.7	768	20.6	53.0	26.4	100	22.3	62.9	14.8	100	10.4	105
North Eastern(Urban)	10.5	83	0.0	62.1	37.9	100	35.1	51.2	13.8	100	0.0	9
Nyanza	19.8	1,161	16.8	67.5	15.7	100	29.6	64.4	6.0	100	6.7	230
Rift valley	14.1	1,775	23.8	53.2	23.0	100	31.4	62.6	6.1	100	10.0	250
Western	20.1	867	14.3	63.9	21.8	100	22.8	69.1	8.1	100	0.3	174
Rural/Urban												
Rural	15.8	5,357	20.9	59.1	20.1	100	29.1	63.8	7.1	100	7.9	847
Urban	16.6	1,909	22.9	66.3	10.7	100	18.7	77.8	3.4	100	4.9	317
Age												
< 6 months	12.4	768	13.1	66.2	20.7	100	17.2	72.8	10.1	100	4.7	95
6-11 months	30.7	782	15.3	68.3	16.4	100	22.2	71.2	6.6	100	3.7	240
12-23 months	22.4	1,544	21.0	61.1	17.9	100	28.7	64.4	6.8	100	6.3	345
24-35 months	14.7	1,470	28.6	57.1	14.3	100	25.6	68.9	5.5	100	9.7	216
36-47 months	10.3	1,430	20.6	55.6	23.8	100	31.4	63.8	4.8	100	7.1	147
48-59 months	9.4	1,272	29.8	56.0	14.2	100	29.4	68.2	2.4	100	13.1	120
Woman's education level												
None	16.3	1,637	14.5	67.6	17.8	100	31.5	64.0	4.6	100	5.3	266
Primary	17.4	3,864	24.7	58.0	17.3	100	27.1	67.4	5.5	100	7.8	673
Secondary +	12.7	1,765	19.9	62.4	17.7	100	17.4	72.9	9.8	100	6.8	224
Total	16.0	7,266	21.4	61.1	17.5	100	26.3	67.7	6.1	100	7.1	1,164

Table 25: Percentage of under-five children with acute respiratory infection in the last two weeks and treatment by health providers, 2000

	Had acute respiratory infection	Number of children under 5	Hospital	Health centre	Dispensary	Village health worker	MCH clinic	Mobile/out reach clinic	Private physician	Traditional healer	Other	Any appropriate provider	Number of children with ARI
Sex													
Male	8.6	3,720	10.6	15.9	13.5	1.3	0.7	0.3	12.3	1.1	19.4	47.7	318
Female	7.7	3,546	11.1	22.9	12.8	3.2	0.4	0.8	7.5	1.6	18.1	52.4	271
Province													
Nairobi	7.2	1,130	9.2	15.9	24.1	4.6	0.0	0.0	4.8	0.0	11.6	58.5	81
Central	7.4	794	13.7	20.3	20.5	2.0	4.5	0.0	15.1	0.0	13.0	56.0	59
Coast	5.7	689	17.7	21.7	15.8	0.0	0.0	0.0	12.1	3.7	20.0	63.5	39
Eastern	6.3	768	14.0	15.6	4.8	0.0	0.0	0.0	15.2	3.6	16.1	49.7	48
North Eastern(Urban	0.3	83	0.0	0.0	0.0	0.0	100.0	0.0	0.0	0.0	0.0	100.0	1
Nyanza	11.6	1,161	14.1	9.7	8.0	4.3	0.4	1.8	9.9	1.5	16.5	42.3	135
Rift valley	6.2	1,775	10.5	21.8	15.8	0.5	0.0	0.6	11.6	0.9	18.0	53.1	109
Western	13.7	867	3.7	29.5	8.2	1.4	0.0	0.0	7.3	1.4	30.8	42.2	119
Rural/Urban													
Rural	8.6	5,357	9.5	20.1	11.9	1.9	0.6	0.6	10.2	1.7	20.8	46.4	461
Urban	6.8	1,909	15.8	15.6	17.9	3.3	0.6	0.3	9.6	0.0	11.7	62.2	129
Age													
< 6 months	5.2	768	17.3	19.6	19.9	2.1	1.4	0.0	21.5	2.0	12.0	66.7	40
6-11 months	8.1	782	11.6	22.5	8.3	0.0	0.0	0.0	15.0	2.6	21.6	56.1	63
12-23 months	10.0	1,544	9.1	22.6	13.8	4.4	0.8	1.6	7.8	2.7	17.9	54.7	154
24-35 months	9.5	1,470	13.3	17.2	12.0	1.3	0.0	0.0	10.9	0.0	14.8	48.9	140
36-47 months	6.3	1,430	9.6	15.7	13.6	2.0	0.0	0.0	8.6	0.8	22.8	44.7	90
48-59 months	8.1	1,272	8.2	17.3	14.1	1.6	1.7	0.6	6.2	0.4	23.0	38.1	103
Woman's education level													
None	7.0	1,637	9.0	17.2	10.9	0.4	0.0	0.5	12.2	1.0	19.2	46.3	115
Primary	9.3	3,864	11.3	20.8	13.5	1.4	1.0	0.7	8.0	1.4	17.8	49.9	359
Secondary +	6.6	1,765	11.3	15.7	14.7	6.4	0.0	0.0	14.5	1.3	21.5	53.5	116
Total	8.1	7,266	10.8	19.1	13.2	2.2	0.6	0.5	10.1	1.3	18.8	49.9	590

World Summit for Children Goal => Number 24

Table 26: Percentage of children 0-59 months of age reported ill during the last two weeks who received increased fluids and continued feeding, Kenya, 2000

	Reported illness in last two weeks	Number of children under 5	Drinking during illness			Total	Eating during illness			Total	Received increased fluids and continued eating	Number of sick children
			More	Same/Less	Missing/DK		somewhat less/same/more	Much less/none	Missing/DK			
Sex												
Male	45.1	3,720	17.9	70.0	12.1	100	31.1	62.9	6.0	100	6.9	1,679
Female	42.1	3,546	17.2	71.3	11.5	100	31.4	61.0	7.6	100	6.5	1,495
Province												
Nairobi	42.8	1,130	13.4	83.9	2.7	100	20.7	78.0	1.4	100	2.5	484
Central	38.4	794	16.6	78.2	5.3	100	32.2	65.2	2.6	100	4.9	305
Coast	52.0	689	31.0	62.5	6.5	100	48.0	50.5	1.5	100	13.0	358
Eastern	39.0	768	18.6	70.4	11.0	100	32.8	60.4	6.7	100	8.8	300
North Eastern(Urban)	19.6	83	0.0	79.8	20.2	100	31.3	61.3	7.3	100	0.0	16
Nyanza	45.2	1,161	18.2	74.7	7.1	100	33.4	63.3	3.3	100	7.4	525
Rift valley	39.1	1,775	18.4	52.6	29.0	100	29.3	51.4	19.4	100	8.8	694
Western	56.8	867	10.6	79.6	9.8	100	28.4	67.3	4.3	100	2.7	493
Rural/Urban												
Rural	43.9	5,357	17.4	68.9	13.7	100	33.0	59.0	8.0	100	7.4	2,349
Urban	43.2	1,909	17.9	75.6	6.5	100	26.3	70.6	3.2	100	4.7	825
Age												
< 6 months	37.8	768	10.2	77.6	12.2	100	23.8	66.5	9.7	100	4.2	290
6-11 months	54.2	782	13.4	72.0	14.6	100	27.3	65.2	7.5	100	4.3	424
12-23 months	49.8	1,544	16.0	71.6	12.4	100	30.2	62.8	7.1	100	6.5	769
24-35 months	46.5	1,470	21.8	69.4	8.8	100	35.8	59.1	5.1	100	9.3	684
36-47 months	36.2	1,430	19.8	64.6	15.6	100	32.3	59.4	8.2	100	5.1	517
48-59 months	38.5	1,272	19.7	71.9	8.4	100	33.3	62.1	4.6	100	8.7	489
Woman's education level												
None	43.0	1,637	14.8	73.7	11.6	100	34.7	60.2	5.2	100	6.7	704
Primary	45.4	3,864	19.1	68.6	12.3	100	31.0	61.9	7.1	100	7.2	1,755
Secondary +	40.5	1,765	16.5	72.6	10.9	100	28.6	64.1	7.3	100	5.4	715
Total	43.7	7,266	17.6	70.6	11.8	100	31.3	62.0	6.7	100	6.7	3,174

Table 27: Percentage of caretakers of children 0-59 months who know at least 2 signs for seeking care immediately, 2000

	Not able to drink/breastfeed	Becomes sicker	Develops a fever	Has fast breathing	Has difficult breathing	Has blood in stool	Is drinking poorly	Child is vomiting	Child has diarrhoea	Knows at least two signs	Number of caretakers
Province											
Nairobi	12.0	22.4	64.7	7.0	7.0	8.7	25.6	35.2	35.0	66.0	864
Central	22.7	25.1	65.3	31.0	29.7	12.5	30.5	41.2	45.3	79.2	597
Coast	19.2	35.5	69.8	19.4	26.6	13.9	18.1	64.9	70.2	84.1	454
Eastern	37.2	32.9	69.6	27.6	32.9	21.2	55.5	57.6	52.8	85.1	560
North Eastern(Urban)	32.4	16.7	72.3	15.5	13.8	8.1	36.1	48.3	65.8	88.5	48
Nyanza	17.8	33.7	55.4	14.1	18.5	5.8	23.7	50.0	51.0	74.7	796
Rift valley	21.3	27.6	55.4	17.4	24.8	8.0	25.5	41.7	46.7	73.8	1,167
Western	21.4	27.5	54.6	25.7	21.6	11.8	31.7	44.0	49.3	77.9	563
Rural/Urban											
Rural	23.2	29.9	58.8	21.4	24.8	11.2	30.8	47.3	50.1	77.7	3,627
Urban	15.2	25.2	66.8	12.2	14.6	9.1	24.4	43.1	45.2	71.8	1,423
Woman's education level											
None	22.1	30.5	56.3	16.8	22.1	10.1	28.3	48.2	52.5	77.9	1,144
Primary	20.7	28.7	60.0	19.0	21.8	9.6	28.3	44.8	46.0	75.1	2,622
Secondary +	20.9	27.0	67.1	21.0	22.6	13.6	31.5	47.4	51.6	76.8	1,283
Total	21.1	28.7	60.9	19.0	22.1	10.7	29.1	46.2	48.8	76.2	5,050

Monitoring IMCI and Malaria Indicator

Table 28: Percentage of children 0-59 months of age who slept under an insecticide-impregnated bednet during the previous night, 2000

	Slept under a bednet		Number of children	Bednet treated		Children who slept under a bednet
	Yes	No		Yes	No	
Sex						
Male	16.2	83.8	3,645	18.9	81.1	609
Female	18.3	81.7	3,471	16.2	83.8	640
Province						
Nairobi	33.7	66.3	1,118	6.1	93.9	379
Central	8.3	91.7	757	7.7	92.3	64
Coast	33.4	66.6	677	17.7	82.3	229
Eastern	16.0	84.0	757	5.8	94.2	121
North Eastern(Urban)	79.5	20.5	80	33.7	66.3	64
Nyanza	18.6	81.4	1,145	39.0	61.0	218
Rift valley	6.1	93.9	1,740	20.0	80.0	109
Western	6.6	93.4	842	23.5	76.5	65
Rural/Urban						
Rural	10.3	89.7	5,235	25.8	74.2	557
Urban	36.5	63.5	1,881	10.9	89.1	692
Age						
< 6 months	22.0	78.0	749	19.3	80.7	167
6-11 months	21.0	79.0	767	18.1	81.9	165
12-23 months	16.2	83.8	1,499	20.7	79.3	247
24-35 months	16.2	83.8	1,446	15.8	84.2	237
36-47 months	16.2	83.8	1,405	16.3	83.7	236
48-59 months	15.6	84.4	1,252	15.1	84.9	197
Total	17.2	82.8	7,116	17.5	82.5	1,249

Monitoring IMCI and Malaria Indicator

Table 29: Percentage of children 0-59 months of age who were ill with fever in the last two weeks who received anti-malarial drugs, Kenya, 2000

	Had a fever in last two weeks	Number of children under 5	Paracetamol	Chloroquine	Fansidar	Other	Don't know	Any appropriate anti-malarial drug	Number of children with fever
Sex									
Male	15.7	3,720	75.1	44.9	25.7	28.6	0.8	65.4	582
Female	15.1	3,546	67.5	40.7	28.9	28.0	0.8	63.3	536
Province									
Nairobi	7.5	1,130	68.3	23.1	27.4	34.2	0.0	50.5	84
Central	6.5	794	66.8	47.4	33.6	15.3	0.0	75.3	52
Coast	18.1	689	78.8	50.1	27.2	21.3	1.5	66.5	125
Eastern	12.6	768	63.5	43.6	11.2	16.1	2.3	53.8	97
North Eastern(Urban)	11.0	83	37.4	39.5	11.1	49.4	0.0	50.6	9
Nyanza	27.2	1,161	74.3	53.0	21.8	26.8	0.2	69.0	316
Rift valley	9.5	1,775	65.2	46.4	30.4	24.8	0.4	69.4	168
Western	30.9	867	74.5	30.6	36.9	39.9	1.2	61.5	268
Rural/Urban									
Rural	17.0	5,357	71.7	44.5	25.8	27.5	0.7	64.6	912
Urban	10.8	1,909	70.5	35.7	33.9	31.8	1.0	63.7	206
Age									
< 6 months	8.4	768	64.8	48.4	19.4	26.4	0.9	60.5	64
6-11 months	16.1	782	71.2	44.1	25.6	36.9	0.0	64.9	126
12-23 months	18.4	1,544	66.5	43.6	27.2	28.0	0.2	64.5	283
24-35 months	17.2	1,470	69.2	39.7	28.3	25.7	0.3	62.5	253
36-47 months	13.7	1,430	80.4	43.6	29.9	31.8	3.1	68.3	196
48-59 months	15.4	1,272	75.0	42.6	27.1	23.7	0.3	63.9	196
Woman's education level									
None	19.4	1,637	65.9	46.9	20.8	23.1	2.0	62.2	317
Primary	15.8	3,864	72.7	43.3	27.4	28.9	0.3	65.4	612
Secondary +	10.7	1,765	76.6	35.0	37.9	34.9	0.0	65.0	189
Total	15.4	7,266	71.5	42.9	27.3	28.3	0.8	64.4	1,118

Table 30: Percentage of women aged 15-49 (Adults) who know the main ways of preventing HIV transmission, 2000

	Heard of AIDS	Percent who know transmission can be prevented by:					Doesn't know any way	Number of women
		Have only one faithful uninfected sex partner	Using a condom every time	Abstaining from sex	Knows all three ways	Knows at least one way		
Province								
Nairobi	96.2	78.6	61.6	78.4	54.2	85.7	14.3	1,656
Central	90.0	81.9	56.2	77.9	52.7	83.4	16.6	1,293
Coast	94.9	78.0	56.1	75.6	52.2	81.6	18.4	909
Eastern	92.0	81.8	46.9	77.6	42.4	84.6	15.4	1,566
North Eastern(Urban)	75.9	53.2	29.0	51.6	28.5	53.7	46.3	98
Nyanza	93.5	77.4	45.7	70.9	36.6	84.5	15.5	1,607
Rift Valley	91.5	79.5	51.7	72.6	45.6	82.6	17.4	2,073
Western	90.0	76.0	49.9	71.7	43.7	81.4	18.6	1,079
Rural/Urban								
Rural	91.9	79.0	49.6	73.6	43.8	82.9	17.1	7,492
Urban	93.9	78.7	59.0	77.6	52.8	84.4	15.6	2,789
Age								
15-19	87.5	70.7	47.3	66.6	40.6	75.7	24.3	2,179
20-24	93.1	80.1	57.8	76.7	51.6	84.9	15.1	2,097
25-29	95.1	83.3	57.5	78.1	50.0	87.8	12.2	1,870
30-34	93.5	80.6	52.6	77.0	47.7	85.2	14.8	1,413
35-39	93.5	80.5	48.4	76.0	43.5	83.6	16.4	1,234
40-44	93.9	82.0	52.5	77.5	47.3	85.9	14.1	927
45-49	93.2	79.2	38.9	73.5	35.7	82.1	17.9	561
Woman's education level								
None	90.1	68.0	39.3	63.9	34.9	71.9	28.1	1,578
Primary	92.4	78.5	50.8	73.3	43.9	83.2	16.8	5,882
Secondary +	93.9	85.9	62.2	83.6	57.3	89.9	10.1	2,821
Total	92.5	78.9	52.2	74.7	46.2	83.3	16.7	10,281

Monitoring HIV/AIDS Indicator

Table 30a: Percentage of women aged 15-19 (Adolscents) who know the main ways of preventing HIV transmission, Kenya, 2000

		Percent who know transmission can be prevented by:							
	Heard of AIDS	Have only one faithful uninfected sex partner	Using a condom every time	Abstaining from sex	Knows all three ways	Knows at least one way	Doesn't know any way	Number of women	
Province									
Nairobi	95.9	72.1	56.1	71.2	49.5	76.8	23.2	335	
Central	80.9	71.1	48.3	66.7	44.2	72.8	27.2	231	
Coast	94.1	75.8	58.5	76.3	52.0	82.4	17.6	167	
Eastern	81.8	64.5	33.4	64.8	27.2	71.7	28.3	341	
North Eastern(Urban)	62.6	44.0	24.7	44.9	24.7	44.9	55.1	26	
Nyanza	91.2	72.0	47.0	63.4	36.2	79.6	20.4	367	
Rift Valley	85.3	70.3	48.9	65.6	42.8	74.2	25.8	422	
Western	86.6	74.1	46.0	65.4	40.6	77.4	22.6	290	
Rural/Urban									
Rural	86.4	71.4	45.5	66.0	38.7	76.3	23.7	1,612	
Urban	90.6	68.6	52.4	68.5	46.1	74.0	26.0	567	
Woman's age									
15	83.3	66.2	44.3	56.9	35.1	70.4	29.6	450	
16	85.3	64.1	41.9	61.0	35.6	70.3	29.7	460	
17	87.3	70.7	46.3	69.8	40.6	76.7	23.3	439	
18	90.0	73.5	50.3	70.4	44.4	78.0	22.0	475	
19	92.5	80.9	55.2	77.3	49.0	84.9	15.1	355	
Woman's education level									
None	82.3	67.9	44.5	62.3	39.3	69.7	30.3	199	
Primary	87.8	68.8	45.3	64.8	38.0	74.7	25.3	1,635	
Secondary +	89.0	80.9	58.2	77.7	53.6	83.7	16.3	345	
Total	87.5	70.7	47.3	66.6	40.6	75.7	24.3	2,179	

Monitoring HIV/AIDS Indicator

Table 30m: Percentage of Men aged 15-54 (Adults) who know the main ways of preventing HIV transmission, 2000

	Percent who know transmission can be prevented by:							Number of Men
	Heard of AIDS	Have only one faithful uninfected sex partner	Using a condom every time	Abstaining from sex	Knows all three ways	Knows at least one way	Doesn't know any way	
Province								
Nairobi	95.5	86.0	62.7	86.7	57.3	91.7	8.3	1,137
Central	99.2	94.5	71.4	88.7	66.7	95.7	4.3	935
Coast	99.5	89.5	71.1	88.7	68.6	93.2	6.8	634
Eastern	99.2	93.2	65.7	87.0	58.6	95.4	4.6	1,139
North Eastern(Urban)	96.3	75.4	38.7	71.5	36.5	76.9	23.1	79
Nyanza	99.4	89.1	62.2	82.2	50.2	95.7	4.3	1,166
Rift Valley	99.1	90.6	67.8	84.6	59.6	94.7	5.3	1,552
Western	98.9	85.6	69.6	81.3	60.4	91.1	8.9	828
Rural/Urban								
Rural	99.2	90.7	66.8	84.5	58.8	94.3	5.7	5,530
Urban	97.0	86.9	65.4	87.4	60.1	92.8	7.2	1,940
Age								
15-19	96.4	80.0	62.6	75.1	52.5	85.6	14.4	1,430
20-24	98.0	87.6	68.5	84.2	58.4	94.4	5.6	1,101
25-29	98.5	91.7	73.8	86.5	65.3	95.7	4.3	1,043
30-34	99.5	93.4	71.4	88.7	65.3	96.5	3.5	1,032
35-39	99.8	94.8	63.2	89.0	57.7	97.3	2.7	899
40-44	99.6	94.8	69.1	91.9	65.6	97.1	2.9	770
45-49	99.3	90.1	62.1	88.6	56.8	96.0	4.0	600
Man's education level								
None	97.9	83.6	59.9	78.5	51.7	88.3	11.7	736
Primary	98.3	88.7	66.2	83.5	58.5	92.7	7.3	4,148
Secondary +	99.4	93.0	68.8	90.0	62.3	97.4	2.6	2,586
Wealth Index Quintiles								
Poorest	98.9	87.2	61.9	81.8	52.5	92.7	7.3	1,449
Second	99.7	90.4	67.2	83.7	59.1	94.2	5.8	1,509
Middle	99.1	90.9	67.9	86.7	61.4	94.6	5.4	1,436
Fourth	97.7	90.6	67.9	84.7	60.8	93.2	6.8	1,526
Richest	97.7	89.5	67.1	89.2	61.7	94.8	5.2	1,550
Total	98.6	89.7	66.4	85.2	59.1	93.9	6.1	7,470

Monitoring HIV/AIDS Indicator

Table 30ma: Percentage of Men aged 15-19 (Adolscents) who know the main ways of preventing HIV transmission, Kenya, 2000

	Percent who know transmission can be prevented by:							Number of women
	Heard of AIDS	Have only one faithful uninfected sex partner	Using a condom every time	Abstaining from sex	Knows all three ways	Knows at least one way	Doesn't know any way	
Province								
Nairobi	77.0	53.9	52.3	54.3	42.3	59.1	40.9	119
Central	98.0	86.2	70.3	78.3	62.1	89.0	11.0	192
Coast	97.8	78.1	67.3	77.7	65.2	83.5	16.5	124
Eastern	98.5	87.9	63.9	78.5	52.3	90.8	9.2	230
North Eastern(Urban)	89.3	57.0	29.6	55.4	29.6	67.8	32.2	9
Nyanza	97.9	79.0	60.4	73.2	43.8	89.4	10.6	260
Rift Valley	97.7	83.1	63.8	81.7	56.7	88.4	11.6	272
Western	99.2	79.9	59.9	73.1	48.8	85.3	14.7	223
Rural/Urban								
Rural	98.2	82.8	63.5	76.4	52.9	88.1	11.9	1,205
Urban	86.6	64.9	57.7	67.7	50.4	72.0	28.0	225
Man's education level								
None	94.2	70.6	67.3	73.5	52.4	81.0	19.0	94
Primary	96.1	79.2	60.1	74.3	50.7	84.6	15.4	1,135
Secondary +	98.8	88.8	74.3	80.1	63.0	93.0	7.0	201
Wealth Index Quintiles								
Poorest	96.9	75.5	58.8	72.3	44.8	84.1	15.9	304
Second	99.3	85.1	62.7	77.6	54.6	89.4	10.6	357
Middle	98.8	83.4	63.8	76.7	53.6	88.2	11.8	280
Fourth	90.8	75.4	61.3	69.9	52.1	79.4	20.6	316
Richest	95.8	80.5	69.5	81.5	60.9	87.0	13.0	173
Total	96.4	80.0	62.6	75.1	52.5	85.6	14.4	1,430

Monitoring HIV/AIDS Indicator

Table 31: Percentage of women aged 15-49 who correctly identify misconceptions about HIV/AIDS, 2000

	AIDS can't be transmitted by			A healthy looking person can be infected	Knows all three misconceptions	Knows at least one misconception	Doesn't correctly identify any misconception	Number of women
	Heard of AIDS	supernatural means	mosquito bites					
Province								
Nairobi	96.2	75.4	62.5	83.2	53.0	90.8	9.2	1,656
Central	90.0	80.9	56.8	78.2	47.3	88.7	11.3	1,293
Coast	94.9	84.2	64.5	73.8	48.2	92.5	7.5	909
Eastern	92.0	83.1	55.6	83.8	48.9	90.2	9.8	1,566
North Eastern(Urban)	75.9	60.1	45.6	42.3	15.4	72.1	27.9	98
Nyanza	93.5	77.3	53.5	79.9	40.5	91.7	8.3	1,607
Rift Valley	91.5	77.0	51.1	72.7	38.5	87.5	12.5	2,073
Western	90.0	76.8	48.6	74.9	36.8	88.1	11.9	1,079
Rural/Urban								
Rural	91.9	79.2	53.3	76.9	41.6	89.4	10.6	7,492
Urban	93.9	77.3	61.8	80.8	51.7	90.1	9.9	2,789
Age								
15-19	87.5	71.3	49.9	69.6	38.9	82.9	17.1	2,179
20-24	93.1	81.1	61.1	79.8	49.5	90.8	9.2	2,097
25-29	95.1	82.4	58.4	83.3	47.5	93.5	6.5	1,870
30-34	93.5	79.9	57.7	81.1	46.0	91.8	8.2	1,413
35-39	93.5	80.8	53.2	79.8	43.3	90.9	9.1	1,234
40-44	93.9	77.3	52.6	76.8	40.0	90.2	9.8	927
45-49	93.2	80.0	52.3	75.6	40.3	89.2	10.8	561
Woman's education level								
None	90.1	73.7	49.3	64.7	31.8	84.7	15.3	1,578
Primary	92.4	78.0	52.6	77.7	41.7	89.5	10.5	5,882
Secondary +	93.9	82.9	65.2	86.0	56.8	92.5	7.5	2,821
Total	92.5	78.6	55.6	78.0	44.3	89.6	10.4	10,281

Monitoring HIV/AIDS Indicator

Table 31m: Percentage of Men aged 15-54 who correctly identify misconceptions about HIV/AIDS, 2000

	AIDS can't be transmitted by:			A healthy looking person can be infected	Knows all three misconceptions	Knows at least one misconception	Doesn't correctly identify any misconception	Number of Men
	Heard of AIDS	supernatural means	mosquito bites					
Province								
Nairobi	95.5	83.9	67.4	89.2	63.6	93.3	6.7	1,137
Central	99.2	85.1	63.5	89.4	57.3	96.4	3.6	935
Coast	99.5	87.1	65.1	87.9	60.0	94.6	5.4	634
Eastern	99.2	87.5	63.1	91.6	57.2	96.7	3.3	1,139
North Eastern(Urban)	96.3	58.3	48.7	76.4	42.7	80.5	19.5	79
Nyanza	99.4	79.3	56.2	86.9	47.1	95.6	4.4	1,166
Rift Valley	99.1	81.8	55.9	84.9	48.7	93.8	6.2	1,552
Western	98.9	82.5	55.6	83.3	45.9	95.2	4.8	828
Rural/Urban								
Rural	99.2	82.7	57.9	86.4	50.3	95.0	5.0	5,530
Urban	97.0	85.0	67.6	90.4	63.2	94.4	5.6	1,940
Age								
15-19	96.4	73.3	48.2	76.7	39.5	88.6	11.4	1,430
20-24	98.0	84.8	63.8	87.7	56.4	94.7	5.3	1,101
25-29	98.5	87.2	63.9	88.2	58.0	96.8	3.2	1,043
30-34	99.5	85.8	65.7	92.1	60.2	96.9	3.1	1,032
35-39	99.8	90.8	65.8	92.9	60.8	98.5	1.5	899
40-44	99.6	86.6	61.0	92.3	55.6	96.9	3.1	770
45-49	99.3	82.7	61.7	87.8	54.8	94.0	6.0	600
50-54	99.7	76.6	57.1	89.2	47.1	96.5	3.5	436
Man's education level								
None	97.9	72.9	51.9	79.8	43.9	89.0	11.0	736
Primary	98.3	81.8	55.1	85.2	48.0	94.0	6.0	4,148
Secondary +	99.4	88.7	71.3	93.2	65.5	97.9	2.1	2,586
Wealth Index Quintiles								
Poorest	98.9	79.9	53.5	82.2	45.4	93.0	7.0	1,449
Second	99.7	83.3	58.8	86.5	51.1	95.2	4.8	1,509
Middle	99.1	84.6	57.1	88.3	51.0	94.7	5.3	1,436
Fourth	97.7	82.7	62.6	89.5	55.7	96.0	4.0	1,526
Richest	97.7	85.8	69.4	90.5	64.5	95.2	4.8	1,550
Total	98.6	83.3	60.4	87.4	53.7	94.9	5.1	7,470

Table 32: Percentage of women aged 15-49 who correctly identify means of HIV transmission from mother to child, 2000

	Percent who know AIDS can be transmitted:				Did not know any specific way	Number of women
	During pregnancy	At Delivery	Through breastmilk	Knows all three ways		
Province						
Nairobi	89.8	73.8	64.9	50.5	6.0	1,656
Central	83.6	76.2	55.8	49.3	12.7	1,293
Coast	85.7	66.1	75.0	56.6	10.2	909
Eastern	83.3	70.7	64.9	52.8	13.6	1,566
North Eastern (Urban)	41.6	37.8	32.6	25.7	53.2	98
Nyanza	82.3	63.6	66.6	49.0	12.7	1,607
Rift Valley	81.8	67.8	60.4	48.1	14.6	2,073
Western	81.3	66.8	59.8	47.6	15.2	1,079
Rural/Urban						
Rural	82.7	68.6	62.8	50.1	13.6	7,492
Urban	85.8	70.2	64.1	49.6	9.7	2,789
Age						
15-19	72.2	55.7	52.1	38.6	23.1	2,179
20-24	85.8	71.3	66.6	52.8	10.3	2,097
25-29	88.7	76.1	64.8	53.1	7.5	1,870
30-34	86.1	70.1	62.2	48.7	10.4	1,413
35-39	85.1	72.1	68.4	54.0	10.7	1,234
40-44	87.0	73.8	68.4	56.8	10.2	927
45-49	86.3	72.0	70.1	55.8	10.4	561
Woman's education level						
None	77.0	61.0	64.7	48.5	19.1	1,578
Primary	82.7	66.9	62.6	48.6	13.2	5,882
Secondary +	88.8	78.1	63.5	53.6	7.7	2,821
Total	83.5	69.1	63.2	50.0	12.6	10,281

Monitoring HIV/AIDS Indicator

Table 32m: Percentage of Men aged 15-54 who correctly identify means of HIV transmission from mother to child, 2000

	Percent who know AIDS can be transmitted:				Did not know any specific way	Number of Men
	During pregnancy	At Delivery	Through breastmilk	Knows all three ways		
Province						
Nairobi	84.7	64.2	56.7	40.6	9.9	1,137
Central	88.0	81.6	58.0	51.3	6.7	935
Coast	90.9	68.5	77.2	58.6	5.8	634
Eastern	86.2	71.5	59.0	44.7	8.7	1,139
North Eastern (Urban)	74.9	65.6	54.8	44.6	21.4	79
Nyanza	85.9	63.6	67.7	47.9	9.1	1,166
Rift Valley	86.8	72.2	63.8	50.4	7.8	1,552
Western	82.7	65.9	58.4	46.3	11.3	828
Rural/Urban						
Rural	86.2	70.9	62.7	49.3	8.7	5,530
Urban	86.1	66.1	61.3	44.1	8.6	1,940
Age						
15-19	72.9	55.1	49.0	35.8	21.0	1,430
20-24	85.9	68.7	62.8	47.3	8.3	1,101
25-29	90.6	69.8	67.4	49.5	5.5	1,043
30-34	89.9	73.9	65.4	49.2	5.4	1,032
35-39	92.4	80.0	67.9	54.4	3.5	899
40-44	91.8	73.9	63.4	53.7	5.2	770
45-49	89.2	73.5	65.4	51.9	4.8	600
50-54	83.8	74.0	65.2	51.6	7.1	436
Man's education level						
None	80.4	65.3	62.8	47.1	13.2	736
Primary	84.4	65.8	61.2	46.9	11.2	4,148
Secondary +	90.7	77.0	64.0	49.9	3.5	2,586
Wealth Index Quintiles						
Poorest	83.7	66.8	64.1	48.8	10.9	1,449
Second	86.4	70.6	62.7	49.9	8.2	1,509
Middle	87.2	71.7	65.2	51.2	8.9	1,436
Fourth	85.4	68.1	57.8	44.2	9.6	1,526
Richest	88.1	71.0	62.0	45.9	6.1	1,550
Total	86.2	69.6	62.3	48.0	8.7	7,470

Monitoring HIV/AIDS Indicator

Table 33: Percentage of women aged 15-49 who express a discriminatory attitude towards people with HIV/AIDS, 2000

	Percent of Women who:				Number of women
	Believe that a teacher with HIV should not be allowed to work	Would not buy food from a person with HIV/AIDS	Agree with at least one discriminatory statement	Agree with neither discriminatory statement	
Province					
Nairobi	49.5	35.1	56.5	43.5	1,656
Central	38.4	32.4	45.8	54.2	1,293
Coast	42.2	38.0	50.3	49.7	909
Eastern	35.2	29.7	41.8	58.2	1,566
North Eastern (Urban)	28.3	22.7	32.5	67.5	98
Nyanza	37.2	38.5	48.6	51.4	1,607
Rift Valley	36.6	26.0	42.8	57.2	2,073
Western	37.9	32.4	47.2	52.8	1,079
Rural/Urban					
Rural	35.7	30.9	43.9	56.1	7,492
Urban	49.0	36.7	55.9	44.1	2,789
Age					
15-19	34.1	27.2	41.5	58.5	2,179
20-24	41.7	34.9	50.1	49.9	2,097
25-29	45.4	36.5	52.5	47.5	1,870
30-34	40.6	33.3	48.1	51.9	1,413
35-39	37.8	32.5	45.8	54.2	1,234
40-44	36.8	32.0	46.9	53.1	927
45-49	34.9	29.3	41.5	58.5	561
Woman's education level					
None	26.9	24.3	33.7	66.3	1,578
Primary	36.3	29.9	44.7	55.3	5,882
Secondary +	52.7	42.5	59.9	40.1	2,821
Total	39.3	32.5	47.2	52.8	10,281

Monitoring HIV/AIDS Indicator

Table 33m: Percentage of Men aged 15-54 who express a discriminatory attitude towards people with HIV/AIDS, 2000

	Percent of Men who:				Number of Men
	Believe that a teacher with HIV should not be allowed to work	Would not buy food from a person with HIV/AIDS	Agree with at least one discriminatory statement	Agree with neither discriminatory statement	
Province					
Nairobi	62.5	40.6	66.3	33.7	1,137
Central	46.3	41.6	53.3	46.7	935
Coast	55.7	44.1	61.5	38.5	634
Eastern	44.8	37.8	52.2	47.8	1,139
North Eastern (Urban)	58.2	56.7	63.3	36.7	79
Nyanza	50.2	45.4	60.7	39.3	1,166
Rift Valley	46.6	37.4	54.2	45.8	1,552
Western	54.8	46.3	63.2	36.8	828
Rural/Urban					
Rural	46.1	39.8	54.4	45.6	5,530
Urban	65.1	46.3	69.7	30.3	1,940
Age					
15-19	40.0	29.5	46.1	53.9	1,430
20-24	51.8	41.1	58.8	41.2	1,101
25-29	55.9	45.0	64.8	35.2	1,043
30-34	55.2	49.0	62.0	38.0	1,032
35-39	56.4	45.5	63.0	37.0	899
40-44	51.6	43.8	60.0	40.0	770
45-49	54.9	40.0	62.3	37.7	600
50-54	47.9	43.4	55.9	44.1	436
Man's education level					
None	43.1	35.8	48.6	51.4	736
Primary	44.3	36.0	52.7	47.3	4,148
Secondary +	64.2	51.9	70.2	29.8	2,586
Wealth Index Quintiles					
Poorest	43.5	39.1	52.4	47.6	1,449
Second	47.9	39.7	55.3	44.7	1,509
Middle	44.3	34.4	52.1	47.9	1,436
Fourth	52.1	43.4	60.3	39.7	1,526
Richest	66.5	50.1	70.8	29.2	1,550
Total	51.1	41.5	58.4	41.6	7,470

Monitoring HIV/AIDS Indicator

Table 34: Percentage of women aged 15-49 who have sufficient knowledge of HIV/AIDS transmission, Kenya, 2000

	Heard of AIDS	Know 2 ways to prevent HIV transmission	Correctly identify 3 misconceptions about HIV transmission	Have sufficient knowledge	Number of women
Province					
Nairobi	96.2	57.1	53.0	35.1	1,656
Central	90.0	55.6	47.3	30.0	1,293
Coast	94.9	54.7	48.2	32.1	909
Eastern	92.0	46.0	48.9	24.8	1,566
North Eastern (Urban)	75.9	28.7	15.4	11.3	98
Nyanza	93.5	41.6	40.5	19.0	1,607
Rift Valley	91.5	50.1	38.5	22.5	2,073
Western	90.0	46.8	36.8	18.9	1,079
Rural/Urban					
Rural	91.9	47.6	41.6	22.3	7,492
Urban	93.9	55.6	51.7	34.6	2,789
Age					
15-19	87.5	44.6	38.9	22.3	2,179
20-24	93.1	55.2	49.5	30.5	2,097
25-29	95.1	54.3	47.5	28.4	1,870
30-34	93.5	50.6	46.0	26.1	1,413
35-39	93.5	46.8	43.3	22.3	1,234
40-44	93.9	50.7	40.0	24.0	927
45-49	93.2	38.0	40.3	20.0	561
Woman's education level					
None	90.1	38.1	31.8	16.2	1,578
Primary	92.4	48.2	41.7	22.3	5,882
Secondary +	93.9	59.8	56.8	38.0	2,821
Total	92.5	49.8	44.3	25.6	10,281

Monitoring HIV/AIDS Indicator

Table 34m: Percentage of Men aged 15-54 who have sufficient knowledge of HIV/AIDS transmission, Kenya, 2000

	Heard of AIDS	Know 2 ways to prevent HIV transmission	Correctly identify 3 misconceptions about HIV transmission	Have sufficient knowledge	Number of Men
Province					
Nairobi	95.5	60.5	63.6	43.7	1,137
Central	99.2	70.7	57.3	44.3	935
Coast	99.5	70.3	60.0	46.4	634
Eastern	99.2	64.0	57.2	41.9	1,139
North Eastern (Urban)	96.3	38.4	42.7	23.4	79
Nyanza	99.4	57.6	47.1	29.9	1,166
Rift Valley	99.1	66.0	48.7	36.0	1,552
Western	98.9	66.2	45.9	33.1	828
Rural/Urban					
Rural	99.2	64.4	50.3	36.2	5,530
Urban	97.0	63.7	63.2	45.4	1,940
Age					
15-19	96.4	59.0	39.5	28.3	1,430
20-24	98.0	65.7	56.4	39.9	1,101
25-29	98.5	71.4	58.0	42.8	1,043
30-34	99.5	69.5	60.2	46.7	1,032
35-39	99.8	61.7	60.8	39.4	899
40-44	99.6	68.0	55.6	41.7	770
45-49	99.3	60.1	54.8	39.0	600
50-54	99.7	57.3	47.1	33.5	436
Man's education level					
None	97.9	57.1	43.9	31.0	736
Primary	98.3	63.7	48.0	34.8	4,148
Secondary +	99.4	67.2	65.5	46.9	2,586
Wealth Index Quintiles					
Poorest	98.9	58.0	45.4	30.7	1,449
Second	99.7	65.0	51.1	37.3	1,509
Middle	99.1	66.0	51.0	38.0	1,436
Fourth	97.7	66.8	55.7	40.9	1,526
Richest	97.7	65.1	64.5	45.6	1,550
Total	98.6	64.2	53.7	38.6	7,470

Monitoring HIV/AIDS Indicator

Table 35: Percentage of women aged 15-49 who know where to get an AIDS test and who have been tested, Kenya, 2000

	Know a place to get tested	Have been tested	If tested, have been told result	Number of women
Province				
Nairobi	74.4	30.9	90.5	1,656
Central	69.9	10.8	59.8	1,293
Coast	66.9	7.3	65.0	909
Eastern	53.6	5.5	78.1	1,566
North Eastern (Urban)	48.0	4.1	100.0	98
Nyanza	61.3	4.1	56.3	1,607
Rift Valley	63.2	6.9	65.1	2,073
Western	57.0	5.2	67.6	1,079
Rural/Urban				
Rural	59.8	5.7	60.9	7,492
Urban	73.9	23.1	88.2	2,789
Age				
15-19	50.3	4.8	71.0	2,179
20-24	65.9	10.9	77.5	2,097
25-29	73.5	14.2	83.9	1,870
30-34	69.6	15.7	74.4	1,413
35-39	65.0	9.7	75.0	1,234
40-44	59.8	10.7	77.2	927
45-49	62.4	5.5	71.1	561
Woman's education level				
None	47.4	6.3	70.7	1,578
Primary	60.8	8.1	69.3	5,882
Secondary +	78.5	17.6	86.3	2,821
Total	63.6	10.4	77.3	10,281

Monitoring HIV/AIDS Indicator

Table 35m: Percentage of Men aged 15-54 who know where to get an AIDS test and who have been tested, Kenya, 2000

	Know a place to get tested	Have been tested	If tested, have been told result	Number of Men
Province				
Nairobi	77.2	17.4	90.3	1,137
Central	77.9	9.4	55.8	935
Coast	82.8	7.7	73.5	634
Eastern	67.9	5.7	82.1	1,139
North Eastern (Urban)	84.1	16.1	100.0	79
Nyanza	78.8	5.9	59.9	1,166
Rift Valley	78.0	5.7	64.9	1,552
Western	72.0	5.6	62.8	828
Rural/Urban				
Rural	73.8	5.7	63.4	5,530
Urban	83.2	15.4	85.8	1,940
Age				
15-19	56.8	2.0	38.2	1,430
20-24	75.7	4.7	62.3	1,101
25-29	80.8	10.2	73.5	1,043
30-34	84.0	12.9	77.6	1,032
35-39	85.7	9.6	72.9	899
40-44	84.5	10.0	76.1	770
45-49	79.4	11.6	89.8	600
50-54	75.5	9.5	85.1	436
Man's education level				
None	64.5	8.7	73.7	736
Primary	72.5	5.2	63.5	4,148
Secondary +	85.6	12.9	81.5	2,586
Wealth Index Quintiles				
Poorest	70.8	4.7	58.4	1,449
Second	73.6	4.4	57.1	1,509
Middle	73.6	5.9	57.6	1,436
Fourth	77.5	8.0	76.1	1,526
Richest	85.2	17.6	86.8	1,550
Total	76.3	8.2	74.3	7,470

Monitoring HIV/AIDS Indicator

Table 37: Percentage of mothers with a birth in the last 12 months protected against neonatal tetanus, 2000

	Received at least 2 doses, last within 3 years	Received at least 3 doses, last within 10 years	Protected against tetanus	Number of mothers
Province				
Nairobi	85.5	0.8	86.3	345
Central	87.4	0.3	87.6	182
Coast	80.7	6.6	87.3	160
Eastern	82.6	0.9	83.6	205
North Eastern (Urban)	100.0	0.0	100.0	19
Nyanza	78.6	1.1	79.7	335
Rift Valley	79.2	1.8	81.0	370
Western	74.7	2.6	77.3	219
Rural/Urban				
Rural	79.4	1.9	81.3	1,324
Urban	86.2	1.3	87.5	511
Woman's education level				
None	79.6	5.0	84.7	235
Primary	78.5	1.7	80.3	1,111
Secondary +	88.3	0.2	88.4	489
Total	81.3	1.7	83.0	1,835

World Summit for Children Goal => Number 22

Table 38: Percent distribution of women aged 15-49 with a birth in the last year by type of personnel delivering antenatal care, 2000

	Person delivering antenatal care							Total	Any skilled personnel	Number of women
	missing	Doctor	Nurse/midwife	Auxiliary midwife	Traditional birth attendant	Other	No antenatal care received			
Province										
Nairobi	2.5	32.4	49.3	0.0	3.0	11.1	1.6	100	81.7	342
Central	0.7	15.7	64.8	3.3	0.0	12.7	2.8	100	83.8	182
Coast	0.7	11.9	54.7	4.4	8.9	13.3	6.1	100	71.0	159
Eastern	0.7	8.8	75.7	0.8	1.6	9.4	3.1	100	85.3	202
North Eastern (Urban)	0.0	4.3	8.4	0.0	0.0	87.4	0.0	100	12.6	19
Nyanza	4.3	5.5	66.8	0.2	9.3	9.1	4.8	100	72.5	335
Rift Valley	2.0	15.7	59.1	1.0	7.1	11.4	3.7	100	75.8	370
Western	0.0	11.3	55.9	1.1	16.9	10.6	4.3	100	68.2	218
Rural/Urban										
Rural	1.9	10.3	62.7	1.5	8.3	11.2	4.2	100	74.4	1,320
Urban	1.9	28.0	52.2	0.4	2.2	13.1	2.2	100	80.7	508
Woman's education level										
None	2.9	6.9	48.1	1.9	11.0	21.5	7.8	100	56.8	234
Primary	2.4	11.5	63.5	1.1	7.5	10.5	3.7	100	76.0	1,105
Secondary +	0.2	27.7	57.1	1.0	2.7	9.8	1.5	100	85.8	489
Total	1.9	15.2	59.8	1.2	6.6	11.7	3.6	100	76.2	1,828

World Summit for Children Goals => Numbers 9, 11

Table 39: Percent distribution of women aged 15-49 with a birth in the last year by type of personnel assisting at delivery, 2000

	Person assisting at delivery						Total	Any skilled personnel	Number of women	
	Doctor	Nurse/midwife	Auxiliary midwife	Traditional birth attendant	Relative/friend	Other/missing				No assistance received
Province										
Nairobi	24.6	39.2	2.6	15.2	5.6	11.9	0.9	100	66.4	342
Central	20.9	41.2	0.0	2.6	16.4	18.9	0.0	100	62.1	182
Coast	9.1	23.2	2.7	24.0	27.2	13.9	0.0	100	34.9	159
Eastern	7.1	38.5	1.2	29.4	12.6	11.2	0.0	100	46.9	202
North Eastern (Urban)	0.0	12.6	0.0	0.0	0.0	87.4	0.0	100	12.6	19
Nyanza	3.8	22.2	0.0	36.7	20.3	16.8	0.3	100	26.0	335
Rift Valley	9.3	23.1	3.3	30.0	20.5	13.6	0.2	100	35.7	370
Western	5.9	23.4	0.8	36.3	17.9	13.4	2.4	100	30.1	218
Rural/Urban										
Rural	7.8	25.6	1.0	29.6	20.3	15.2	0.5	100	34.4	1,320
Urban	21.4	39.3	3.1	14.9	6.5	14.2	0.6	100	63.8	508
Woman's education level										
None	7.4	16.7	0.8	29.2	20.1	24.4	1.3	100	25.0	234
Primary	7.5	26.1	1.2	30.2	20.0	14.8	0.3	100	34.8	1,105
Secondary +	22.7	42.9	2.8	13.4	6.7	10.7	0.8	100	68.4	489
Total	11.6	29.4	1.6	25.6	16.4	14.9	0.5	100	42.6	1,828

World Summit for Children Goal => Numbers 11

Table 40: Percent distribution of children aged 0-59 months by whether birth is registered and reasons for non-registration, 2000

	Registration status								Reason DK or Missing	Birth certificate issued	Number of children
	Birth registered	DK if birth registered	Costs too much	Must travel too far	Didn't know it should be registered	Late & didn't want to pay fine	Doesn't know where to register	Other			
Sex											
Male	62.4	1.6	1.3	4.3	14.7	3.7	4.4	2.8	4.8	12.3	3,720
Female	63.1	1.4	1.5	3.6	15.9	4.0	3.2	2.5	4.8	14.7	3,546
Province											
Nairobi	82.9	2.1	0.3	1.5	6.8	1.1	1.2	0.9	3.2	24.1	1,130
Central	90.7	0.3	0.6	1.1	3.4	0.6	1.1	0.7	1.5	16.0	794
Coast	64.1	0.7	1.2	5.1	16.2	3.0	2.3	3.0	4.4	13.4	689
Eastern	64.2	0.9	1.4	3.3	15.6	4.8	4.3	1.5	3.9	8.5	768
North Eastern'	88.0	0.0	0.0	2.5	0.9	4.6	0.0	1.4	2.4	72.2	83
Nyanza	41.5	1.6	2.7	7.1	25.0	7.2	6.8	2.5	5.4	10.0	1,161
Rift valley	52.1	1.3	1.7	5.2	20.7	3.8	5.2	3.6	6.4	9.9	1,775
Western	56.5	3.1	1.4	3.0	13.3	5.8	3.6	6.1	7.4	8.2	867
Rural/Urban											
Rural	56.1	1.5	1.6	4.9	17.9	4.7	4.7	3.2	5.5	8.5	5,357
Urban	81.6	1.5	0.7	1.5	7.8	1.4	1.3	1.1	3.0	27.4	1,909
Age											
< 6 months	58.9	0.9	1.7	4.1	16.0	3.4	4.2	5.9	4.9	5.6	768
6-11 months	63.7	1.5	1.7	3.9	16.4	3.0	3.0	2.5	4.3	9.8	782
12-23 months	61.9	1.1	1.4	3.8	15.7	3.8	4.7	2.0	5.5	11.2	1,544
24-35 months	64.8	1.4	1.0	4.1	14.0	4.9	2.8	3.3	3.7	14.6	1,470
36-47 months	63.4	2.5	1.3	3.9	15.2	3.9	3.6	1.6	4.6	17.4	1,430
48-59 months	62.5	1.1	1.6	4.0	15.1	3.5	4.3	2.2	5.7	17.5	1,272
Woman's education level											
None	52.2	2.4	2.1	5.7	20.8	3.8	4.7	3.3	5.0	10.1	1,637
Primary	58.1	1.3	1.5	4.2	17.2	4.5	4.7	3.0	5.6	9.6	3,864
Secondary +	82.8	0.9	0.7	1.9	5.8	2.5	1.0	1.3	3.0	25.0	1,765
Total	62.8	1.5	1.4	4.0	15.3	3.8	3.8	2.7	4.8	13.5	7,266

Table 41: Percentage of children 0-14 years of age in households not living with a biological parent, 2000

	Living with neither parent					Living with mother only		Living with father only		Impossible to determine/ Missing	Total	Not living with a biological parent	One or both parents dead	Number of children
	Living with both parents	only father alive	only mother alive	both are alive	both are dead	father alive	father dead	mother alive	mother dead					
Sex														
Male	66.5	0.5	0.5	3.2	1.3	16.5	4.7	1.5	0.7	4.6	100	5.6	7.8	10,929
Female	64.9	0.8	0.9	3.9	0.9	16.4	5.3	1.2	0.7	4.9	100	6.5	8.7	10,846
Province														
Nairobi	73.3	0.4	0.6	0.9	1.1	12.2	2.0	0.5	0.1	8.8	100	2.9	4.2	2,392
Central	62.1	0.5	0.3	2.8	0.8	18.4	4.0	0.8	0.2	10.1	100	4.4	5.8	2,427
Coast	65.9	0.9	0.8	4.0	0.4	17.6	3.5	2.1	0.9	3.9	100	6.1	6.5	1,926
Eastern	62.7	0.4	0.2	3.1	0.5	19.6	6.5	2.2	0.7	3.9	100	4.3	8.3	3,482
North Eastern (Urban)	74.1	2.0	2.3	5.0	1.9	7.9	3.4	0.0	0.4	3.1	100	11.2	10.0	196
Nyanza	63.1	1.1	1.2	2.9	3.1	15.4	7.4	0.6	1.1	4.1	100	8.3	13.9	3,653
Rift valley	70.0	0.2	0.6	3.8	0.6	15.7	4.9	1.2	0.6	2.4	100	5.2	6.9	5,081
Western	60.7	1.3	1.3	7.3	1.1	16.7	5.0	2.1	1.3	3.2	100	10.9	10.0	2,617
Rural/Urban														
Rural	64.1	0.7	0.8	3.9	1.2	17.2	5.6	1.3	0.8	4.4	100	6.5	9.1	17,439
Urban	72.0	0.5	0.6	2.0	1.0	13.5	2.6	1.3	0.3	6.3	100	4.1	5.0	4,336
Age														
0-4 years	69.9	0.4	0.2	1.7	0.3	18.5	2.9	0.5	0.3	5.1	100	2.6	4.2	8,247
5-9 years	65.5	0.7	0.8	4.3	1.1	15.5	5.2	1.6	0.7	4.6	100	6.9	8.5	6,877
10-14 years	60.7	0.9	1.3	5.0	2.2	14.7	7.4	2.0	1.2	4.5	100	9.4	13.0	6,651
Total	65.7	0.6	0.7	3.6	1.1	16.4	5.0	1.3	0.7	4.7	100	6.0	8.3	21,775

Monitoring Children's Rights Indicator

Table 42: Percentage of children 5-17 years of age who are currently working, Kenya, 2000

	Paid work	Unpaid work	Domestic work		working on Own/family farm/business	Currently working	Number of children
			< 4 hours/day	4 or more hours/day			
Sex							
Male	2.6	2.9	52.9	4.5	37.1	9.0	8,216
Female	3.1	2.6	60.6	8.3	31.3	12.6	8,325
Province							
Nairobi	4.2	0.2	33.2	4.3	0.6	8.5	1,481
Central	3.2	2.6	64.6	2.7	35.4	7.4	1,951
Coast	4.0	2.0	46.3	7.6	29.6	12.7	1,430
Eastern	3.2	3.0	60.2	9.1	33.5	14.0	2,822
North Eastern (Urban)	5.0	0.0	32.4	9.5	14.8	12.5	131
Nyanza	1.9	3.5	69.5	4.4	41.6	8.5	2,794
Rift valley	2.0	2.4	51.0	6.9	35.9	10.2	3,801
Western	2.9	4.6	63.8	8.4	48.6	14.0	2,131
Rural/Urban							
Rural	2.6	3.2	60.8	6.6	40.4	11.1	13,678
Urban	4.0	0.5	37.6	5.6	4.5	9.5	2,863
Age							
5-9 years	0.7	0.4	48.4	4.0	21.0	5.0	6,877
10-14 years	2.8	1.8	66.5	6.3	42.0	10.1	6,651
15-17 years	7.6	10.3	54.6	12.2	47.0	25.8	3,013
Total	2.8	2.8	56.8	6.4	34.2	10.8	16,541

Monitoring Children's Rights Indicator

Table 43: Percentage of under-five children who are severely or moderately undernourished, 2000

	Weight for age: -2 SD		Weight for age: -3 SD		Height for age: -2 SD		Height for age: -3 SD		Weight for height: -2 SD		Weight for height: -3 SD		Number of children	
	MICS 2000	KDHS 1998	MICS 2000	KDHS 1998	MICS 2000	KDHS 1998	MICS 2000	KDHS 1998	MICS 2000	KDHS 1998	MICS 2000	KDHS 1998	MICS 2000	KDHS 1998
Sex														
Male	22.6	22.2	6.7	4.6	37.9	35.2	15.7	13.2	6.6	5.9	1.8	1.5	3,000	2246
Female	19.6	22.0	4.6	5.0	32.6	30.8	13.7	12.1	5.3	6.2	0.9	1.4	2,917	2167
Province														
Nairobi	12.4	11.4	1.8	2.1	29.6	25.7	11.1	7.1	3.1	7.1	0.8	3.6	760	257
Central	15.4	14.3	3.3	2.0	27.4	27.5	9.6	9.8	4.6	5.6	1.1	1.7	627	414
Coast	21.1	27.4	4.0	7.1	33.7	39.1	15.5	18.3	6.4	4.3	1.5	1.0	566	346
Eastern	29.6	25.7	6.9	6.6	42.8	36.8	17.4	13.2	7.8	4.7	1.4	0.9	673	753
North Eastern(Urban)	16.6		3.7		35.4		17.0		7.2		0.0		65	
Nyanza	19.9	22.2	6.2	5.8	35.9	30.8	16.3	11.2	5.2	7.0	1.7	1.2	959	906
Rift valley	24.9	24.9	7.3	4.2	36.8	33.1	14.9	13.3	7.6	7.4	1.5	1.8	1,533	1134
Western	21.5	19.1	8.0	3.6	38.1	35.0	16.9	14.3	5.5	4.6	1.4	0.9	734	604
Rural/Urban														
Rural	23.9	23.9	6.8	5.3	38.0	34.7	16.0	13.8	6.8	6.2	1.5	1.3	4,497	3662
Urban	12.4	13.3	2.1	2.3	26.6	24.7	10.4	7.4	3.3	5.1	0.8	2.0	1,420	751
Age														
< 6 months	3.0	2.3	0.5	0.4	12.4	7.1	1.4	0.8	2.4	5.2	0.8	1.3	322	427
6-11 months	14.6	14.8	4.2	4.6	24.5	17.5	6.2	4.8	3.9	7.8	0.6	2.5	598	506
12-23 months	28.4	26.9	9.2	7.2	47.5	41.8	21.3	17.5	9.9	9.1	2.5	1.6	1,312	979
24-35 months	22.5	28.3	6.6	6.3	34.8	37.8	13.5	13.4	6.6	4.9	1.4	0.9	1,290	909
36-47 months	19.4	23.8	3.8	3.3	34.5	35.6	15.3	15.3	4.1	4.9	1.0	1.8	1,286	811
48-59 months	21.9	22.7	4.9	4.0	34.7	38.0	16.1	14.7	4.9	4.0	0.9	0.8	1,109	781
Woman's education level														
None	24.1	36.8	7.2	8.8	37.2	46.4	17.2	21.4	7.1	8.8	1.6	1.2	1,347	487
Primary	23.3	{26.9,19.9"}	6.8	{6.1,4.2"}	38.8	{39.7, 31.5"}	16.3	{16.4,11.3"}	6.6	{6.5,6.4"}	1.7	{1.6,1.7"}	3,147	{1657,1118"}
Secondary +	13.7	11.0	1.8	1.7	25.6	19.2	8.9	5.0	3.4	3.9	0.5	1.0	1,423	1152
Total	21.2	22.1	5.7	4.8	35.3	33.0	14.7	12.7	6.0	6.1	1.4	1.4	5,917	4413

World Summit for Children Goal => Number 3, 9

Table 44: Percentage of children age 12-23 months currently vaccinated against childhood diseases, 2000

	BCG		DPT 1		DPT 2		DPT 3		Polio 0	Polio 1		Polio 2		Polio 3		Measles		All		None		with health		of children			
	MICS 2000	KDHS 1998		MICS 2000	KDHS 1998	MICS 2000	KDHS 1998	MICS 2000	KDHS 1998	MICS 2000	KDHS 1998																
Sex																											
Male	90.3	95.0	88.1	95.5	82.6	90.6	77.3	80.5	79.8	84.7	95.4	79.9	91.4	73.1	81.8	76.5	79.4	61.9	66.0	6.7	2.4	64.3	55.9	770	549		
Female	91.5	96.7	90.8	96.0	82.8	89.3	75.1	77.8	81.6	88.5	95.3	82.3	89.5	72.5	79.7	75.8	79.1	59.7	64.8	4.7	2.9	66.4	55.0	774	548		
Province																											
Nairobi	95.5	97.7	92.4	95.5	86.4	90.9	84.8	75.0	93.9	93.9	97.7	85.0	88.6	74.3	86.4	84.9	93.2	64.0	72.7	3.0	2.3	61.1	40.9	225	81		
Central	88.8	100.0	87.1	100.0	84.5	98.6	82.7	97.2	81.7	83.3	97.8	81.1	95.6	77.6	94.9	80.2	91.3	69.0	84.8	5.9	0.0	64.9	54.8	147	81		
Coast	93.7	94.8	91.9	97.9	81.3	89.5	78.2	81.3	84.1	91.2	98.4	84.6	94.2	75.1	85.3	76.8	91.6	61.1	71.9	4.2	0.8	74.7	65.8	150	100		
Eastern	91.5	97.9	90.5	97.4	83.9	93.6	76.0	86.2	80.0	86.9	96.3	79.9	93.7	74.5	88.9	83.2	86.6	68.2	74.9	6.0	2.1	63.6	69.3	173	173		
North Eastern	95.7		90.0		78.5		72.8		90.0	90.0		84.2		84.2		90.0		72.8		4.3		74.2		21			
Nyanza	85.7	92.9	83.3	92.1	72.0	81.6	61.7	66.2	72.6	75.7	91.2	69.4	79.8	56.9	66.4	68.4	61.9	47.7	46.5	10.2	5.2	57.2	47.1	257	224		
Rift valley	93.2	96.3	91.7	96.5	87.4	92.2	80.7	83.9	86.4	89.7	96.2	86.8	95.1	80.3	84.4	79.5	83.7	68.2	69.3	4.5	2.4	70.4	51.2	378	290		
Western	86.3	94.4	88.6	94.4	82.7	89.2	70.6	72.2	61.2	84.7	94.4	79.3	89.2	69.2	72.0	58.1	65.8	45.4	56.2	6.2	2.8	65.2	61.4	191	148		
Rural/Urban																											
Rural	89.3	95.3	88.3	95.4	81.9	89.9	74.1	79.2	76.5	84.0	95.0	79.3	90.2	71.5	79.9	73.1	76.1	58.7	64.2	6.7	2.9	65.1	58.5	1,150	888		
Urban	95.6	98.0	92.8	97.1	85.1	90.5	82.4	79.0	93.1	94.2	96.8	86.4	91.5	76.3	84.2	85.0	92.8	66.9	70.5	2.9	1.6	66.1	42.5	394	210		
Woman's education level																											
None	85.1	88.1	82.2	89.8	73.2	78.7	66.0	68.1	71.1	80.0	88.8	71.8	81.2	63.3	68.2	68.5	70.1	53.5	53.4	10.7	7.4	59.8	51.4	318	117		
Primary	91.7	95.3	90.3	94.0	82.8	87.8	75.6	75.2	80.0	86.2	97.4	81.4	93.4	72.2	83.3	74.8	82.2	58.2	67.1	4.9	3.5	66.1	57.5	851	13,291		
Secondary +	94.0	98.7	93.6	98.2	90.5	95.1	86.2	86.5	90.6	92.9	97.7	88.3	96.1	82.1	90.2	85.7	89.7	72.8	79.2	3.4	0.6	68.4	53.9	375	276		
Total	90.9	95.9	89.4	95.8	82.7	90.0	76.2	79.2	80.7	86.6	95.4	81.1	90.4	72.8	80.8	76.1	79.2	60.8	65.4	5.7	2.7	65.4	55.4	1,544	1097		

Table 45: Percentage of children 12-23 months immunized against childhood diseases at any time before the survey, Kenya, 2000

		BCG	DPT1	DPT2	DPT3	Polio 1	Polio 2	polio3	Measles	All	None	Number of children
Vaccinated at any time before the survey												
According to:												
Vaccination Card	MICS 2000	63.2	63.3	61.0	58.4	62.5	61.0	57.7	51.2	47.8	0.2	
	KDHS 1998	54.7	54.8	53.9	50.9	54.9	53.8	51.0	46.0	44.2	0.0	
Mother's Report	MICS 2000	27.7	26.1	21.7	17.8	24.1	20.1	15.1	24.9	13.0	5.6	
	KDHS 1998	41.2	41.0	36.1	28.2	40.5	36.6	29.7	33.3	21.2	2.7	
Either	MICS 2000	90.9	89.4	82.7	76.2	86.6	81.1	72.8	76.1	60.8	5.7	1,544
	KDHS 1998	95.9	95.8	90.0	79.1	95.4	90.4	80.7	79.3	65.4	2.7	1,097
Vaccinated by 12 months of age												
	MICS 2000	90.5	88.9	82.2	75.1	85.7	80.3	71.7	71.6	55.6	0.0	1,544
	KDHS 1998	94.0	94.5	88.2	76.0	94.2	89.0	77.7	70.7	58.0	4.4	1,097

World Summit for Children Goal => Number 22

Table 46: Percentage of live births in the last 12 months that weighed below 2500 grams at birth, Kenya, 2000

	Percent of live births below 2500 grams		Number of live births	
	MICS 2000	KDHS 1998	MICS 2000	KDHS 1998
Province				
Nairobi	9.9	5.9	342	219
Central	8.4	8.6	182	306
Coast	9.2	6.1	159	284
Eastern	10.0	4.1	202	584
North Eastern(Urban)	2.3		19	
Nyanza	8.9	1.7	335	753
Rift Valley	9.4	3.4	370	867
Western	9.4	1.6	218	451
Rural/Urban				
Rural	9.2	3.2	1,320	2828
Urban	9.5	6.4	508	636
Woman's education level				
None	8.4	3.2	234	373
Primary	9.5	{3.5,2.8"}	1,105	{1348,870"}
Secondary +	9.3	5.3	489	873
Total	9.3	3.8	1,828	3464

World Summit for Children Goal => Number 12

"Primary complete

Table 47: Birth weight and size at birth

	Birth weight							
	<2500		2500+		DK/Missing		Number	
	MICS 2000	KDHS 1998	MICS 2000	KDHS 1998	MICS 2000	KDHS 1998	MICS 2000	KDHS 1998
Province								
Nairobi	2.8	5.9	55.0	73.1	42.2	21	304	219
Central	3.2	8.6	39.3	71.0	57.5	20.4	159	306
Coast	3.8	6.1	29.8	34.1	66.5	59.9	140	284
Eastern	8.6	4.1	38.0	45.1	53.4	50.8	184	584
North Eastern(Urban)	0.0		0.0		100.0		2	
Nyanza	2.5	1.7	18.0	34.8	79.4	63.5	306	753
Rift Valley	3.0	3.4	25.2	37.4	71.8	59.2	332	867
Western	1.6	1.6	19.7	23.7	78.6	74.7	193	451
Rural/Urban								
Rural	3.3	3.2	24.7	36.2	72.0	60.7	1,178	2828
Urban	3.7	6.4	50.2	64.1	46.1	29.5	441	636
Woman's education level								
None	3.3	3.2	19.9	24.0	76.8	72.8	184	373
Primary	2.9	{3.5, 2.8"}	26.8	{26.2, 45.8"}	70.3	{69.9, 51.4"}	994	{1348, 870"}
Secondary +	4.6	5.3	48.6	66.9	46.8	27.8	442	873
Total	3.4	3.8	31.8	41.3	64.8	54.9	1,620	3464

"Primary complete

Table 48: Percentage of women aged 15-49 (Adults) who know the main ways of preventing HIV transmission, 2000

Percent who know transmission can be prevented by:								
	Using a condom every time		Abstaining from sex		Doesn't know any way		Number of women	
	MICS 2000	KDHS 1998	MICS 2000	KDHS 1998	MICS 2000	KDHS 1998	MICS 2000	KDHS 1998
Province								
Nairobi	61.6	56.9	78.4	35.9	14.3	12.4	1,656	768
Central	56.2	31.4	77.9	17.5	16.6	19	1,293	830
Coast	56.1	38.1	75.6	14.3	18.4	23.9	909	597
Eastern	46.9	32.5	77.6	43.4	15.4	17.4	1,566	1378
North Eastern(Urban)	29.0		51.6		46.3		98	
Nyanza	45.7	38.0	70.9	33.1	15.5	12.9	1,607	1687
Rift Valley	51.7	35.2	72.6	22.6	17.4	20.8	2,073	1648
Western	49.9	37.4	71.7	16.0	18.6	24.4	1,079	896
Rural/Urban								
Rural	49.6	33.1	73.6	26.7	17.1	19.9	7,492	5983
Urban	59.0	52.3	77.6	31.9	15.6	12.2	2,789	1821
Age								
15-19	47.3	33.2	66.6	33.4	24.3	24.9	2,179	1827
20-24	57.8	44.2	76.7	26.4	15.1	14.8	2,097	1537
25-29	57.5	45.0	78.1	24.9	12.2	14.1	1,870	1356
30-34	52.6	{36.4}	77.0	{26.8}	14.8	{16.5}	1,413	{1962}
35-39	48.4	{36.4}	76.0	{26.8}	16.4	{16.5}	1,234	{1962}
40-44	52.5	{28.5}	77.5	{26.8}	14.1	{19.5}	927	{1122}
45-49	38.9	{28.5}	73.5	{26.8}	17.9	{19.5}	561	{1122}
Woman's education level								
None	39.3	17.9	63.9	20.1	28.1	30.9	1,578	860
Primary	50.8	{31.8, 40.8"}	73.3	{26.5, 27.3"}	16.8	{24.7, 16.3"}	5,882	{2871, 1773"}
Secondary +	62.2	49.5	83.6	33.2	10.1	6.6	2,821	2300
Total	52.2	37.5	74.7	27.9	16.7	8.1	10,281	7803

Monitoring HIV/AIDS Indicator